

BIOLOGICAL INVENTORY OF THE RÍO AROS, SONORA, MEXICO: A RIVER UNKNOWN

Final Report to T&E, Inc.

20 June 2006

Chris O'Brien[†], Aaron D. Flesch¹, Eric Wallace², Michael Bogan³, Samia E. Carrillo-Percástegui⁴,
Sky Jacobs and Charles van Riper III



Red-spotted toad (*Bufo punctatus*) Photo: Michael Bogan.

†Corresponding Author: Chris O'Brien
obrienc@email.arizona.edu
University of Arizona and
Sonoran Desert Research Station
Biological Sciences East, 125.
University of Arizona
Tucson, AZ 85721
(520) 623-3720
obrienc@email.arizona.edu

¹ Contributed flora and bird sections. Email: flesch@ag.arizona.edu

² Contributed reptile and amphibian section. Email: batrachia@yahoo.com

³ Contributed aquatic invertebrate section. Email: bogan@lifesci.ucsb.edu

⁴ Contributed mammal section. Email: samiac@email.arizona.edu

CONTENTS

ABSTRACT.....	2
INTRODUCTION.....	3
MATERIALS AND METHODS.....	4
Vegetation and Flora	5
Aquatic Invertebrates	5
Herpetofauna.....	7
Birds.....	8
Mammals.....	8
RESULTS AND DISCUSSION.....	10
Vegetation and Flora	10
Aquatic Invertebrates.....	15
Herpetofauna	18
Birds	23
Mammals	28
CONCLUSIONS	30
ACKNOWLEDGEMENTS.....	31
LITERATURE CITED	31
FIGURES.....	36
TABLES	39
PHOTOS.....	55

ABSTRACT

The area surrounding the Río Aros canyon and upper Río Yaqui valley constitute one of northwestern Mexico's largest and least fragmented wildlands, and is near the boundaries of subtropical and temperate biogeographic regions. Because of its remote and wild character, biogeographic setting, and increasing interest in the region by conservationists and naturalists, we surveyed the river corridor and major tributaries to document biological diversity in the region. In July and August of 2005, we traversed 184.8 km, of the Ríos Aros and Yaqui in inflatable boats from Nátorá to El Río, conducting a preliminary biological inventory of plants, aquatic invertebrates, amphibians and reptiles, birds, and mammals along the river corridor and in tributary canyons. We report range extensions for species of plants (*Ficus pertusa*, *Sideroxylon persimile*, *Tabebuia impetiginosa*), aquatic invertebrates (*Buenoa albida*, *B. thomasi*, *Macrovatellus mexicana*, *Martarega mexicana*), herpetofauna (*Drymarchon corais*, *Terrepene nelsoni*) and observations of birds at the edge of their ranges (*Ara militaris*, *Haliaeetus leucogaster*, *Icterus wagler*, *Vireo flavoviridis*). In total we observed 102 species of aquatic invertebrates, 8 amphibians, 13 reptiles, 80 birds, and 18 mammals during an initial assessment of the biological diversity of these taxa in the region. We hope that this preliminary assessment will aid in future conservation and management efforts in the Río Aros region.

INTRODUCTION

The Río Yaqui basin is the largest watershed in northwestern Mexico and covers 79,162 km², or 30% of the land area of the state of Sonora (Revenga *et al.* 1998). This watershed delivers 70% of the surface flow for northwestern Mexico (Gallo-Reynoso *et al.* 2002), and is a conservation area of concern by the Mexican Government (Arriaga *et al.* 2002a, 2002b). The Río Yaqui drainage contains large tracts of roadless areas that have been little explored by naturalists, but it also serves as an important economic area for forestry, agriculture, ranching, and mining, in northwestern Mexico (Bojorquez *et al.* 1985).

The Río Yaqui's largest tributary, the Río Aros, is a major river that drains much of northeastern Sonora and western Chihuahua, and is at the biogeographic confluence of both subtropical (Sinaloan) and temperate (Madrean, Sonoran, and Chihuahuan) flora and fauna. This region is extremely rugged, accessible by only foot, horseback, or boat, and harbors one of the largest unfragmented wild areas of foothills thornscrub in the state of Sonora (Lorenzana-Piña *et al.* 2004). It supports the largest known population of jaguars (*Panthera onca*) in northern Mexico (Brown and Lopez-González 2001), which, in 2003, prompted the purchase of the 10,000 acre Los Pavos-Northern Jaguar Preserve to serve as a core area for their protection. This reserve, purchased by Naturalia and operated in cooperation with the Northern Jaguar Project and Defenders of Wildlife, has fostered increasing biological and conservation interest in the area.

Recent surveys within the Yaqui basin have noted new, northernmost populations of several species of birds, including military macaw (*Ara militaris*), yellow-green vireo (*Vireo flavoviridis*), and fan-tailed warbler (*Euthlypis lachrymose*) that occur in isolated patches of tropical deciduous and tropical semideciduous evergreen forest and thornscrub (Russell and Monson 1998, A. Flesch, unpubl. data). On the Aros in our study area, tropical river otter (*Lontra longicaudis annectens*) (B. Brown, pers. comm.) and spotted box turtle (*Terrapene nelsoni*; S. Carrillo-Percástegui, unpubl. data) have also been recently observed.

The Yaqui basin is also at the southern range boundary for temperate species such as bald eagle (*Haliaeetus leucocephalus*) (Brown and Warren 1985), beaver (*Castor canadensis frondator*) (Gallo-

Reynosa *et al.* 2002), and is at the southwestern limits of the ranges of lowland leopard frog (*Rana yavapaiensis*) (Platz and Frost 1984) and Chiricahua leopard frog (*R. chiricahuensis*) (Platz and Mecham 1979). Because of this biogeographic setting combined with a lack of basic biological information for the region, we predicted that biological surveys would yield observations of species outside of their currently known geographic ranges. To test this prediction we conducted a baseline biological inventory along the Aros and upper Yaqui river corridor and in select tributaries in the summer of 2005.

METHODS AND MATERIALS

We traversed the Río Aros and upper Yaqui from Nátora (700 m) to El Río (350 m) in inflatable boats during the 2005 monsoon period to ensure adequate flows for river travel (Figs 1, 2). We adopted an expeditionary approach using inflatable kayaks and oar-power rafts, and transported all supplies and equipment for the duration of the trip. Rafting logistics were coordinated by Caiman Expeditions (Lane Larson, Tucson AZ) who provided knowledge of the local area and rafting equipment. Local support in Sahuaripa provided transport to the launching point of Nátora located 100 km east over two-track, unimproved roads.

From Nátora, the Aros flows north, through alternating steep rolling hills and narrow barrancas, passing around the Sierra los Pavos and joining the Río Bavispe to form the Río Yaqui. Below the confluence, the Yaqui flows south towards Presa El Novillo (Fig. 2). We ended our trip at El Río.

For the duration of the study, floating surveys were conducted on an opportunistic basis, with occasional stops in areas of interest. We rafted during the day, traveling between 3-7 km/hr and averaged 26 river km per day, stopping to camp at areas that afforded access to major tributary canyons that we surveyed the following morning (Table 1). We selected tributaries for surveys based on their size and aspect because our primary objective was to document vegetation communities and associated fauna that were rare in the regional landscape or at the edges of their geographic ranges. Tributaries were surveyed for 2-3 hours, starting at dawn, and sometimes additionally for 1-2.5 hours in the early evening. We hiked tributary canyons, surveying the plants, aquatic invertebrates, herpetofauna, birds, and large

mammals using the following methods. Additionally we also visited a high-elevation site en route to Nátora (Figure 2).

Vegetation and Flora

We qualitatively described the flora of the study area, identifying many species of trees and shrubs while focusing on uncommon and rare species of distributional or conservation interest. We noted the distribution and approximate abundance of species of interest and identified plants with the use of botanical keys and specimen vouchers located at the University of Arizona Herbarium or by using our knowledge of woody plants of the region. Many of our observations along the river corridor were from boats; binoculars allowed identification of many tree species growing on canyon walls or steep slopes. We occasionally landed a boat to collect plants of interest along the river, and when on the ground, we focused our efforts in tributaries and near campsites. Because space with which to protect and transport plants was limited, we only collected species of interest and often used digital photographs to aid identification.

Aquatic Invertebrates

We employed a variety of survey techniques to collect aquatic insects in the mainstem river and tributaries. We vigorously swept pools in flowing (Photo 1a) and non-flowing side canyons with a 1-mm mesh d-frame net (d-net) until we visually detected no new species. We sampled riffles in flowing side canyons by disturbing a 0.1m² patch of substrate to a depth of 5 cm, and caught insects downstream with a 1-mm and/or a 325- μ m mesh d-net. High flows and sediment loads made it impossible to take benthic kick samples in the mainstem (Photo 1b), so instead surface netting and drift samples were employed to collect specimens. We collected surface-dwelling and free swimming insects from the upper 0.5 m of the water column in the main river via vigorous sweep netting with a 1-mm mesh d-net. Drift samples were collected between dusk and dawn as drift rates are elevated during this time (Williams and Hynes, 1976). We placed a 325- μ m mesh d-net on the bottom substrate, ensuring that the upper portion of the net was submerged and held in this position for 10 min for each sampling event. High turbidity and large

amounts of detritus prevented longer drift sampling times. All pool, benthic and drift samples were preserved in 95% ethanol and transported to Oregon State University for identification.

Many aquatic insects have extended aerial stages as adults during the summer months, often as a strategy to avoid flash floods (Lytle 2003). Therefore, we visually surveyed both the mainstem and tributary canyons during daylight hours, and employed blacklights after dark on the mainstem. We employed blacklights, both by placing the light against a white sheet hung in riparian areas adjacent to the river, and by placing a clear container filled with ethanol over a light at the river's edge; in all cases lights were deployed for 90 min. We preserved all insects in 70% ethanol and transported them to Oregon State University for identification (SEMERNAT Permiso #SGPA/DGVS/04147).

At most sites where aquatic insects were collected, we recorded water temperature, pH, and conductivity using a hand-held multi-meter. We recorded sampling locations with a Garmin Etrex Vista GPS unit. Flows in tributary canyons were visually estimated, and flows in the mainstem were visually estimated in relation to the known flow amount at the gauging station near Rancho Los Pavos (see Table 1).

We used non-metric multidimensional scaling (NMS), to compare the community structure of different sites sampled within the Río Aros basin. NMS, an ordination technique, reduces redundancy in the data matrix, based on a rank-based metric to reduce community data from many dimensions to a two or three dimensional ordination, and does not make distributional assumptions of the data matrix (McCune and Grace 2002). Environmental parameters were then overlain on the community composition-based ordination to elucidate relationships between measured parameters and community changes along the gradient. Sorensen distances were used to calculate community dissimilarity between samples and regions. Since collection techniques varied between samples, and therefore absolute numbers of individuals varied greatly between samples, abundance data were presence-absence transformed before any ordination analyses. We used Multi-response Permutation Procedure (MRPP) with Sorensen distances to quantify and test within-group agreement and distinctness (see Mielke and Berry 2001) between mainstem river and tributary canyon samples. This procedure yields two statistics,

an A-statistic ($-1 \leq A \leq 1$), describing the effect-size of the grouping, and a p -value, which evaluates the likelihood that observed differences are due to chance. We performed all community analysis using the program PC-ORD, (McCune and Mefford 1999).

Herpetofauna.

We surveyed 14 localities for herpetofauna (Table 4), focusing on major tributary canyons of the Ríos Aros and Yaqui and made incidental observations adjacent to the river and at campsites. We emphasized our efforts on riparian and aquatic environments as we expected a greater diversity of species to be found in these habitats. In addition we recorded observations of easily observed herpetofauna in the uplands en route to and from survey canyons.

We surveyed for aquatic and semi-aquatic herpetofauna using visual encounter surveys (VES), usually during daylight hours and often during early morning (Crump and Scott 1994). We silently approached aquatic environments to visually search and listen for active or escaping herpetofauna. We searched along the water's edge and used d-nets and flashlights to probe vegetation and undercut banks for hidden herpetofauna. Once a perimeter search was complete, we waded through pools to access undercut banks and boulder piles unapproachable from the perimeter. Following VES, we dipnetted runs, riffles and pools for amphibian larvae and aquatic turtles or snakes. VES and dipnetting efforts were more intensive around perennial pools. Digital photographs were taken of most pools and aquatic environments. We also searched under cover objects (e.g. logs, small boulders, debris piles) to detect species likely to be inactive during our surveys. Additionally, we listened for breeding vocalizations of frogs and toads.

Encountered herpetofauna were recorded based on visual observation if identification was certain, or captured to confirm positive identification. For target species captured, we recorded sex and pertinent morphological measurements (e.g. snout-vent length (SVL), color pattern descriptions). We documented locations of target species observations with hand-held GPS units and photographed captured individuals using digital cameras. Based on previous observations from the area (S. Carrillo, unpublished data), target species included ranid frogs (*Rana pipiens* complex, *R. tarahumarae*), indigo snakes (*Drymarchon*

corais), and spotted box turtles (*Terrepenne nelsoni*). Photographs provided a permanent record of our observations and will allow herpetologists familiar with the species of the region to confirm questionable identifications (e.g. leopard frogs). We accessioned photographs into the University of Arizona Herpetological Collections.

Birds.

We used two approaches to examine the distribution and abundance of birds, one along the main river and another along its tributaries. While on or near the river we noted presence of all bird species detected either aurally or visually and recorded number of individuals, pairs, or dependant young for species that were less common. Due to the challenges of simultaneously navigating the river, surveying, and data recording, we only estimated abundance of species of interest; our knowledge of bird vocalizations facilitated identification of nearly all audible species. We surveyed eight major tributaries or side canyons for 1 to 5 km and recorded the numbers of individuals, pairs, flocks, and singing males of each species. We often mimicked a pygmy-owl call to increase detections (Marshall 1957). Along each river segment and tributary we noted evidence of breeding and recorded nest contents when possible. We noted the start and end points of each survey and estimated the length of river segments and tributary transects using 1:50,000-m scale topographic maps. We also described the structure and composition of vegetation at sites where we observed bird species of interest. To compare bird communities among or between river segments and tributary transects we calculated species richness and frequency of occurrence (no. detected/transects) of each species. To quantify and describe breeding and residency status we used the criteria of the Arizona Breeding Bird Atlas (Corman and Wise-Gervais 2005), published information (e.g. Russell and Monson 1998), and our own knowledge and observations. All tributary surveys were completed within approximately 5 hrs after sunrise or initiated within 2 hrs of sunset. Surveys along the river corridor were conducted throughout the day.

Mammals.

We employed four main techniques to document the mammal fauna of the Río Aros region: camera traps, mechanical live-traps, track and scat searches, and opportunistic sightings. We used a 35

mm camera trap (Deercam Inc.) to increase detection rates of secretive or nocturnal species. These cameras, which are triggered by a passive infrared sensor that detects heat and movement (Lynam 2002), were set opportunistically along trails where tracks and minimal cattle activity were detected. Our camera was left mounted to a tree trunk overnight and retrieved the following morning.

To sample terrestrial species of small mammals, we used 20 Sherman traps that were set overnight on a 400 m transect oriented along trails at the mouths of tributary canyons or along the Aros mainstem. Traps were baited with oatmeal and coarsely ground peanut butter. Animals captured were identified, photographed, and released. All traps were set in foothills thornscrub or broadleaf riparian forest.

We searched for tracks and scat opportunistically on beaches along the river corridor during midday stops and at evening camps. Additionally, tributary canyons (Table 1) were systematically searched for evidence of large mammals. We especially focused on areas of soft wet sand and wet or dry mud where tracks were more likely to be preserved. Easily identified tracks were recorded and a photo was taken when possible. Large felid tracks were traced on a piece of Plexiglas with a non-permanent marker and then traced again on a plastic bag with a permanent marker to copy the exact track for future identification confirmation. Measurements of tracks were taken following the protocol established by Childs (1998). We also identified scrapes that large cats make on trunks to sharpen their claws and mark their territories. Scats were identified based on appropriate measurements, diet contents, shape, and association with distinctive tracks (Elbroch 2003). Large felid scats (jaguar and mountain lion) were collected and stored in a paper bag with GPS location and observations and deposited at the University of Arizona for further DNA analysis as part of an ongoing Master's thesis on large cats in the region (S. Carrillo-Percástegui, unpubl. data). All smaller scats were left on site and documented with photographs and location.

Finally, while floating the river and during stops and tributary surveys, we opportunistically recorded all visual sightings of large mammals. When possible, photo vouchers were taken to confirm identification, and location and vegetation types were recorded.

RESULTS AND DISCUSSION

Between 27 July and 4 August 2005 we traversed 184.8 km of the Ríos Aros and Yaqui, surveying river reaches, areas around campsites, and 26.5 km of tributaries. We camped at seven sites along the river and surveyed 11 tributaries of the Río Aros. Figure 1 provides a map of the region, the study area is shown in Figure 2, and all survey sites are shown in Table 1.

At the water levels we encountered of 56 to 114 m³/s (1977-4025 feet³/sec) the Aros is largely a class III river of moderate difficulty, requiring constant scouting and maneuvering. The river was easily negotiated at monsoon discharges, except for one class IV rapid above Carrizoso canyon where the river cut through a more resistant layer of limestone. The river level fluctuated daily as the result of large monsoon storms in local areas and far upstream. For example, on 30 July, between 06:30 and 10:00, flow in the mainstem increased such that the water level increased 1.2 m in vertical distance

Land use along the river consisted primarily of grazing, and we found evidence of cows in nearly all river-side camps and in many of the surveyed tributaries. Additionally, there were stock trails leading away from the river at important access points and at many of the larger tributary mouths. Several small dwellings were observed that are likely used seasonally or intermittently. Results of surveys by taxonomic group are outlined below.

Vegetation and Flora

The Aros and Yaqui rivers flow through the lower western foothills of the Sierra Madre Occidental. Vegetation in this region is comprised of tropical-subtropical scrublands (Brown 1982), specifically foothills thornscrub that grows along the eastern edge of the Sonoran desert and at higher elevations within the Sonoran Desert (Felger *et al.* 2001).

Upland vegetation was dominated by foothills thornscrub; *Lysiloma microphyllum* and *L. watsonii*, *Bursera fagaroides*, *B. laxiflora*, *Ceiba acuminata*, *Jatropha cordata*, *Fouquieria macdougalii*, *Acacia cochliacantha*, *Ipomoea arborescens*, and *Stenocereus thurberi* were widespread and common. We observed palms (*Brahea brandegeei* and *Sabal uresana*) in scattered locations often in small stands

on slopes, and they were common only in a few canyons such as Los Alisos and El Carrizoso. Flowering *Hintonia latifolia* and *Parthenium tomentosum*, were less common and found scattered on slopes in thornscrub. Hecho cactus (*Pachycereus pecten-aboriginum*), was uncommon along most of the Aros corridor and locally common along the Yaqui, often on low slopes and flats near the river. Oaks were restricted to upper elevations often on north-facing slopes along the upper Aros corridor. Small stands of *Quercus chihuahuensis* occurred on hillsides and in drainages along some tributaries often >1 km from the river. We observed *Q. tuberculata* mixed with thornscrub species in uplands and as a riparian species in the first 40 river km below Nátora above 680 m in elevation; *Q. tuberculata* was also a dominant riparian tree above 600 m in the north-facing Canyon Los Lobos (Fig. 1). We observed many large areas of upland thornscrub, often >5 ha in size, on slopes near the bottom of the river canyon or on north-facing slopes where recent frosts had killed most vegetation to the ground (Photo 5). Most of these areas were in the first 60 river km below Nátora and none were observed along the Yaqui. In many cases, new branches and foliage of uniform height had re-sprouted from the base of the top-killed woody trees, most commonly *Acacia cochliacantha*, and was approximately 1/6 to 1/3 the height of standing dead material. The relative short height of dead vegetation, which rarely exceeded approximately 5 m, suggested a recurring history of frost damage in these areas.

We observed associations of several species of riparian trees along the river. Narrow bands of broadleaf riparian woodland dominated by *Salix gooddingii* and *S. bonplandiana* were common on shallow sandy terraces. Other riparian woodlands, dominated by *Prosopis velutina*, *Havardia mexicana*, *Sapindus saponaria*, and *Vitex mollis*, occurred on sandy to rocky terraces that also often harbored *Celtis reticulata*, *Ceiba acuminata*, and *Lysiloma watsonii* above the high-water line; where steep slopes extended down to the river's edge these woodlands occupied narrow linear bands one tree wide. We found no riparian vegetation in the rockiest and steepest areas. *Cephalanthus salicifolius* was fairly common along the river's edge generally well below the high-water line and had large flood-damaged trunks anchored to rock outcrops or depressions. *Guazuma ulimifolia* was common along the Yaqui and in some tributaries but only at lower elevations at and below Los Alisos Canyon (Fig. 1). *Sideroxylon*

persimile was locally common in riparian vegetation in the upper portion of the Aros river corridor and not found above 650 m. We first observed this broadleaf semideciduous tree approximately 12 river-km below Tunapa and at only 1 location on the Yaqui just above the Sahuaripa bridge. The largest and tallest stands of *S. persimile* were in Canyon Los Lobos, where it reached heights of >15 m and dominated the canopy along with *Plantanus racemosa* that was rare along the upper river corridor. *P. racemosa* was only common in Canyon Los Lobos, rare in Canyon El Placer, and seen at the mouths of several smaller side canyons from the river.

In addition to *S. persimile*, we observed only two other species of tropical broadleaf semideciduous trees: *Ficus pertusa* was found in moist situations and *F. petiolaris* was common on rock walls and piles along the river corridor and in sheltered canyons. The uppermost occurrences of *F. pertusa* along the Aros were several trees approximately seven to eight meters tall in a deep, sheltered glen where the river dropped through a narrow canyon approximately three km below the mouth of Canyon Los Lobos at approximately 500m elevation. We also observed *F. pertusa* at three locations along the Río Yaqui within 10 river km of the confluence with the Río Aros; additionally, a small flood-scoured plant was found along a deep rocky canyon draining the Sierra Los Pavos northeast of the confluence with the Aros. Aside from the mixed broadleaf-deciduous (e.g. *Platanus*) and broadleaf semideciduous (e.g. *S. persimile*) riparian vegetation in Canyon Los Lobos, we did not observe other riparian vegetation dominated by tropical semideciduous species.

We observed other species of interest, including several *Tabebuia impetiginosa* trees that were roughly 5 to 6 m tall in El Placer Canyon. *Bernardia cinerea*, a euphorbiaceous shrub or short tree with tomentose leaves was in flower and uncommon, occurring mainly in riparian areas below 500 m. Other plants we identified that are of interest include the woody shrubs *Sebastiania bilocularis*, *Karwinskia humboltiana*, *Zanthoxylum fagara*, *Croton sonore*, *Esenbeckia hartmanii*, *Tecoma stans*, *Guaiacum coulteri*, *Euphorbia colletioides*, and *Randia sonorensis*, vines such as *Funastrum clausum*, grasses such as *Lasiacis ruscifolia*, and the trees *Cordia sonore*, *Eysenhardtia orthocarpa*, *Alvaradoa amorphoides*, and *Jatropha cordata*. We did not observe other species more typical of tropical deciduous forest and

more southern thornscrub such as *Brongniartia alamosana*, *Bursera penicillata*, *Lonchocarpus hermannii*, or *Wimmeria mexicana*.

Conclusions. Much of the vegetation we observed was typical in both structure and composition of that found in the northern portion of the thornscrub, and many species more typical of tropical deciduous forest and southern thornscrub were absent (Brown 1982, Gentry 1982, Martin *et al.* 1998, Robichaux and Yetman 2000). Although the structure and composition of thornscrub along the Río Aros seemed similar to that further south, compared to the surrounding landscape, thornscrub in the Aros Valley is somewhat isolated and connected to more extensive areas of thornscrub only by a narrow low-elevation corridor created by the river canyon. In fact, thornscrub in this area, which continues on up the river corridor as an ever-narrowing band of subtropical lowlands into neighboring Chihuahua, is at the north-easternmost extent of its distribution (Brown and Lowe 1980) and is nearly surrounded by Madrean oak, pine, and mixed conifer forest and grassland.

Although we observed several species associated with tropical deciduous forest or more southern thornscrub to the south including *B. cinerea*, *S. persimile*, and *F.s pertusa*, lower minimum temperature and lower precipitation likely limits the distribution of many species that are found to the south and west. Inversions of cold air during winter resulting from steep regional gradients in elevation may prevent other more tropical species from occupying lowlands. The Aros canyon is close to the highlands of the Sierra Madre Occidental of nearby Chihuahua that reach a maximum of 2,820 m elevation within 60 km of our study site and the mountains east of Bacadehuachi that reach 2,560 m elevation within 50 km. Just 10 km northeast of the Río Aros at Tunapa, the Mesa Cienega de San Jorge reaches an elevation of 1,920 m. As a result, freezing air from these highlands may occasionally penetrate some lowland canyons thereby limiting the distribution of frost-sensitive tropical species. In the case of plant distribution, it is often the extreme, not mean, environmental conditions that are limiting and even an occasional severe frost every few decades may have major impacts on plant distribution (Taylor 1934, MacArthur 1972). Our observations of large patches of frost-killed thornscrub suggest freezing events occur regularly, especially in the upper portion of the Aros corridor.

Frost is possibly more important than precipitation in limiting plant distribution and community structure along the Aros than it is to the south and west of our study area. For example, along the lower Río Bavispe to the west where maximum elevations above the river canyon are lower and adjacent areas of highland much smaller, at least one canyon supports a large riparian gallery forest of the tropical semideciduous species *F. pertusa* and *F. petiolaris* at similar latitude and elevation as those we visited along the Aros (A. Flesch unpubl. data). Still further to the west in the neighboring Moctezuma Valley, where adjacent highlands are much lower and smaller, more diverse tropical riparian associations of *Ficus trigonata*, *F. pertusa*, *Sideroxylon persimile*, *Drypetes gentryi*, and *Xylosma flexuosum* occur in a wet canyon at approximately 700 m, again at similar latitudes to those we visited (A. Flesch unpubl. data). In comparison to these areas, riparian vegetation was never dominated by broadleaf semideciduous elements anywhere along the Aros except to some degree in Canyon Los Lobos. Riparian areas with broadleaf semideciduous structure and physiognomy are characteristic of Sinaloan riparian evergreen forest (Minckley and Brown 1982a). These areas, especially when found north of 29° north latitude, are of great conservation interest as they often harbor isolated and mesic refugia that support the northernmost populations of many plant and animal species typically found only far to the south.

As with other taxa, we observed plants species of distributional interest with nearest known localities that were far to the south. Species of note include *S. persimile* and *F. pertusa* that were approximately 100 km north of their known ranges (Felger *et al.* 2001); the latter species occurs still further to the north in canyons draining into the Río Bavispe near 29° 30' north latitude (A. Flesch unpubl data). The nearest published locality for *Tabebuia impetiginosa* is also approximately 100 km south (Turner *et al.* 1995, Felger *et al.* 2001); this species has also been recorded further to the north in canyons draining into the Río Bavispe (A. Flesch unpubl data). Observations of *Cephalanthus salicifolius* are among the most north-easternmost (Felger *et al.* 2001). This tree has a limited distribution in Sonora and considered a variety of the more widespread *C. occidentalis* (T. van Devender pers. comm.). We are unsure of the significance of *Euphorbia colletioides*, *Funastrum clausum*, and *Bernardia cinerea* in the region.

Time available to study the flora was limited due to expedition time constraints and the complexity of river and overland navigation. Despite these challenges we gained a firm overview of the floristic composition and distribution of vegetation communities in a remote region that has not been previously described.

Aquatic Invertebrates

We detected at least 102 aquatic insect taxa from the Ríos Aros and Yaqui and tributaries during our surveys (Table 3), representing eight orders and 42 families. The vast majority of taxa we collected were aquatic beetles (Coleoptera) and true bugs (Hemiptera). Species richness at individual collection sites varied from two species (Río Aros at Río Bonito) to 47 species (Carrizoso Canyon). Average number of species observed per site was 17. Side canyon tributaries supported 93% of all species we detected (95 species). In contrast, the mainstem river supported only 19 species. Seven of the mainstem species, found in slower edgewater and eddies or in drift, were exclusive to the mainstem, and not observed in tributaries.

Community composition varied significantly by habitat type. NMS ordination of the community samples converged on a stable, and significant solution (stress = 15.5; final instability = 0.001; $p = 0.0099$) illustrating that tributary and mainstem environments supported different communities with no overlap in ordination space (Figure 3, $A=0.15$; $p=0.0001$ for MRPP analyses). Community composition in tributaries varied little among sites especially when compared to variation between mainstem and tributary samples. When physical environmental characteristics (temperature, pH, and conductivity) were overlain on the NMS ordination, only flow characteristics described community composition (Axis 1; $r=0.95$). Clearly, the disparity between tributary flows (0-3 cfs) and mainstem flows (1000-3500cfs) was very large, so the very strong association between flow and community composition is not surprising. The range of temperature (20-29°C), pH (7.25-8), and conductivity (60-590 μ S) may not have been large enough to detect a significant effect on community composition. Alternatively, a lack of data from some sites (Table 2) may have made it difficult to detect the effects of these parameters on community composition.

We collected six aquatic insect species in drift samples, but in extremely low numbers. Flow dynamics in the mainstem likely contributed to the very low numbers of individuals in the drift. Unfortunately, all mainstem drift samples were collected after a large increase in flow, so the drift densities were likely diluted by increased flow. In the Río Aros, only several individuals of three taxa were collected (riffle beetles: Elmidae; midges: Chironomidae; net-spinning caddisflies: Hydropsychidae). The drift collection site on the Río Yaqui yielded four taxa (blackflies: Simuliidae; mayflies: Leptophlebiidae; moths: Pyralidae; net-spinning caddisflies: Hydropsychidae), and higher overall densities than sites on the Aros. The Río Yaqui was much less turbid than the Aros, and appeared to have not been as affected by the increased flows given its wider channel, perhaps explaining the higher drift rates in the Yaqui. Only one tributary, El Placer, had sufficient flow to collect drift and kick samples; despite five replicate samples not one individual insect was collected. This tributary is likely normally dry and recent rainfall had increased flow at the site, resulting in the lack of sufficient time for the recolonization of newly-wetted habitats.

Light trapping on three different nights yielded numerous terrestrial and three aquatic insect species. The giant water bug (*Abedus vicinus sonorensis*) was collected at lights on two nights. This collection is significant because nearly all *Abedus* species are flightless, and examples of *A. v. sonorensis* flying to light are rare in the literature. Adults of the dobsonfly *Corydalus* were abundant at light traps (Photo 4a). *Corydalus* larvae (Photo 4b) were also present in large numbers in the riparian areas immediately adjacent to mainstem; the larvae of this genus are normally completely aquatic, but high humidity and fluctuating river levels may have lead to larval individuals foraging out of the water as well. On the lower Río Aros and on the Río Yaqui, large numbers of the leptophlebiid mayfly *Traverella* were collected at lights. Incredible numbers of *Traverella* individuals emerged from the mainstem Rios Aros and Yaqui at both dawn and dusk; subimago males were abundant adjacent to the river, while mature adults were present in large swarms of thousands of flying individuals, and also congregated on mesquite trees at the river's edge.

The aquatic insect fauna of the Río Aros and its tributaries is very similar to that of streams in the Madrean Sky Islands (MSI) to the north and west of the Aros basin. Nearly 87% of the species collected in the Aros basin are also present in MSI streams (for description of MSI stream fauna see Bogan 2005). Of the 13% of species that did not overlap, many are species with more subtropical distributions (e.g. *Rhantus calidus*, *Buenoa thomasi*, *Abedus breviceps*). In Sonora, these species have typically been found in lower latitude streams, such as the Río Cuchujaqui in southern Sonora (M. Bogan, unpublished data). Thus, the Río Aros represents a northern and upper elevation outpost in the distributions of many subtropical aquatic insect taxa.

This study produced range extensions for several aquatic insect species. The dytiscid beetle *Macrovatellus mexicana* has been observed in isolated ranges southwest of the Aros basin (Bogan 2005: Sierra Aguaje near Guaymas, Sonora) and from the east of the Sierra Madre Occidental (Jasper and Challet, 2004), but not in between. Collections at Carrizoso and Lobos Canyons act to fill a large gap in the distribution of this beetle. The backswimmer bug *Martarega mexicana* was historically known only from central Mexico southward. In the 1960's it was first collected in the Salt-Gila River Basin of Arizona (Truxal and Menke 1966), leaving a distributional gap of nearly 1,500km. In 2004 this species was found in the Cajon Bonito, an extreme northern tributary of the Río Yaqui (Bogan, 2005), and that collection, together with its presence in the Río Aros and Río Yaqui, indicates that the distribution of this species is not as disjunct as previously thought. Instead, more surveys are needed to fully understand the distribution of *M. mexicana* between these two distributional extremes. Two other species of backswimmers collected in the Río Aros, *Buenoa albida* and *B. thomasi*, represent northern extensions of their previously known ranges. The previous northernmost collections of these species were from the Río Mayo in southern Sonora (Truxal 1953).

Conclusions. The Río Aros and its tributaries support diverse aquatic insect communities. Community structure varied dramatically between the mainstem river and tributary canyons. Tributary canyons supported most of the diversity during our summer sampling period. This pattern is likely due to the relative flow stability of tributary canyons during the monsoon season and the relative ease of

sampling these habitats thoroughly. However, the tributary aquatic insect diversity recorded in this study is likely a modest estimate. At many tributary sites we only sampled for as little as 30 min due to time constraints and were still detecting new species when we departed. With increased sampling time we would surely have detected many more species. Aquatic insect diversities from the mainstem habitats are certainly dramatic underestimates, because while monsoon discharge in the Río Aros is conducive to boating, these flows make aquatic insect sampling difficult. There is likely a relatively diverse benthic fauna in the mainstem that was undetected during our survey due to the high, fluctuating flows and very high turbidity. Future studies in the Río Aros basin should focus on more time-intensive benthic surveys of tributary habitats and winter, spring, or fall surveys would allow better assessment of river-dwelling species. The position of the Río Aros as a transitional basin between the Madrean Sky Islands and the subtropical portion of the Sierra Madre Occidental is clear from the aquatic insect communities present in the basin. This transitional location, combined with the additional aquatic insect diversity yet to be described in the basin, make the Río Aros basin an important area for future surveys, exploration, and conservation.

Herpetofauna

Amphibians. We observed eight species of amphibians including one salamander (*Ambystoma rosaceum*) that was seen on the access road between Sahuaripa and Nátorá, and seven species of frogs and toads representing five families (Table 5). We recorded three species of true toads (Bufonidae), two species of true frogs (Ranidae) and one species each of narrow-mouthed toads (Microhylidae) and treefrogs (Hylidae).

Tarahumara salamander (*Ambystoma rosaceum*) larvae were found in a plunge pool at the high elevation site in Madrean evergreen woodland vegetation (Figure 1, Table 4), approximately 10 km SW of the Río Aros. Mazatlán toads (*Bufo mazatlanensis*) were found in the upper reaches of the study area at Nátorá and Tunapa, and Buena Vista and were active and abundant around campsites, likely due to the warm, humid conditions. Sonoran desert and red spotted toads (*B. alvarius* *B. punctatus*) were each found at three sites in low numbers. We encountered small *Bufo* tadpoles at several sites but positive

identification at this life stage is very difficult and so were not included in Table 4.

Leopard frogs (*Rana* cf. *yavapaiensis*) were by far the most common amphibian encountered (n=6 sites, Table 4). We heard what seemed to be a leopard frog vocalizing in Chino Gordo at the only deep plunge pool in the canyon, yet further intensive VES and dip netting failed to confirm presence. We observed leopard frogs most often in low numbers ($n \leq 3$) in and around perennial pools. Most individuals encountered were juveniles. Leopard frogs were found to be relatively abundant at Los Lobos where we observed individuals of all life stages except eggs (i.e., tadpoles, juveniles and adults). We captured four large, robust adults at the only large perennial pool surveyed in the canyon. Females (n=3) measured 72-74 mm SVL and the male measured 61.5 mm SVL. We captured one juvenile Tarahumara frog (*R. tarahumarae*) in La Ciénega in a small cobble and boulder stream. Most ranids captured had light infestations of trombiculid mites (*Hannemania* spp.) on the posterior thigh and ventral surfaces.

We recorded the diminutive and highly secretive Sinaloan narrow-mouthed toad (*Gastrophryne olivacea mazatlanensis*) at La Ciénega and Carrizoso. The former observation was a gravid female (34 mm SVL) in a moist, cobbled wash with no standing water and the latter was a single tadpole from a long, muddy pool lined with dense common reed (*Phragmites communis*). We encountered tadpoles and adults of canyon treefrogs (*Hyla arenicolor*) in four rocky canyon sites and heard one individual calling at Carrizoso (Table 4).

Noteworthy amphibian observations include those of leopard frogs (*Rana* cf. *yavapaiensis*), and Tarahumara frog (*R. tarahumarae*). We tentatively assigned leopard frogs encountered to *R. yavapaiensis*, a conservative approach because leopard frogs are notoriously morphologically conserved making field identification of closely related species difficult. The Northwest Mexico leopard frog (*R. magnaocularis*) is sister species to *R. yavapaiensis* (Hillis and Wilcox 2005) and little definitive morphological data exists with which to distinguish these 2 species in the field (Frost and Bagnara 1976, Platz and Frost 1984). The extent to which lowland leopard frogs range south into Mexico is unknown (Platz and Frost 1984) and the northern extent of the Northwest Mexico leopard frog (*R. magnaocularis*) is also little known (Frost and Bagnara 1976). Both species are found in similar aquatic environments

from near sea level to 1500m.

The northern most published record for Northwest Mexico leopard frog is ~40 km south of our study area although this species assignment is based on one individual and a morphological characteristic (incompleteness of supralabial stripe) that is common to both species (Lemos-Espinal *et al.* 2004b). Based on our experience with *R. yavapaiensis* and published accounts of Northwest Mexico leopard frogs we could find no obvious deviations from typical *R. yavapaiensis* from Arizona and therefore tentatively assign our observations to the latter species. Considerable work needs to be done to better delineate these 2 species' ranges and, their potential zone of sympatry, and also to develop diagnostic field characters for field identification.

Leopard frogs have suffered precipitous declines throughout the southwestern United States (Clarkson and Rorabaugh 1989) including *R. yavapaiensis*. The extent to which these declines may be occurring in northwestern Mexico are unknown primarily due to a lack of baseline distributional data on leopard frogs and the general inaccessibility of this region. Most recently, a newly described pathogenic chytrid fungus implicated in declines of amphibians worldwide, including pristine wilderness areas, has been identified in leopard frogs in Arizona (Bradley *et al.* 2002). Leopard frogs were relatively common in our area and are known to be so on the Los Pavos Northern Jaguar Preserve (P. Warshall pers. comm.) suggesting that populations in the area are wide-spread. Certainly, more extensive (e.g. inventory) and intensive (e.g. chytrid sampling) work needs be done in order to provide a baseline of species presence and their distribution, abundance, and population status in order to develop conservation and management strategies for leopard frogs in the region.

Tarahumara frogs are relatively well documented from the sierras of northeastern Sonora and were considered to be experiencing population declines in parts of this region during the 1980's and 1990's (Hale *et al.* 1998). We observed only one juvenile frog in La Ciénega Canyon, suggesting that it was a dispersing individual from a population higher in the drainage. The area surveyed in La Cienega contained very few rocky plunge-pools that are considered habitat for Tarahumara frogs (Stebbins 2003). Based on our observation, other populations may exist in the basin that could be important when

considering the conservation status of this species.

Reptiles. We observed 12 species of reptiles representing seven families, including three species of turtles, seven species of lizards, and 3 species of snakes. Turtles represented two families (Emydidae, Kinosternidae), lizards four families (Helodermatidae, Phrynosomatidae, Scincidae, Teiidae) and snakes one family (Colubridae) (Table 5).

We found one slider shell (*Trachemys scripta* subsp.), a largely riverine group, on the beach at Buena Vista and made several observations of basking individuals as we floated the river. There is only one slider native to the Yaqui River basin (*Trachemys scripta yaquia*) (Ernst and Barbour 1989) but there have been reports of an introduced subspecies (*Trachemys scripta* subsp.) in the basin (C. Schwalbe pers. comm.). Despite the remote nature of our study area we would be remiss to assign our observation to a particular subspecies based solely on a weather-beaten shell and visual observations made from a distance. The most commonly encountered turtle was the Sonoran mud turtle (*Kinosternon sonoriense*); we observed both adults and young of the year hatchlings. We recorded mud turtles from five localities, four of which were rocky canyons, their typical aquatic habitat (Table 4).

We observed spotted box turtles, a terrestrial species of interest, at Buena Vista camp and in Los Lobos canyon. The Buena Vista individual, found in a dry arroyo with foothills thornscrub appeared to be a female based on plastron shape and length of tail (Ernst and Barbour 1989). The Los Lobos individual was also found in a dry arroyo in foothills thornscrub just above a perennial reach. This individual had a crimson stained beak, suggesting it had been eating organ-pipe cactus fruit in the area.

Tree lizards (*Urosaurus ornatus*) and Clark's spiny lizards (*Sceloporous clarkii*) were common arboreal species found throughout the study area and were often observed on trees and rocks around camp and in tributary canyons. Whiptails (*Aspidoscelis* spp.) were the most abundant surface dwelling species encountered, although none were captured and no positive identifications to species were made. We observed one Great Plains skink (*Eumeces obsoletus*) at the gauging station below Buena Vista. No photos were taken before this skink escaped into a deep rock crack above the river. A Gila monster (*Heloderma suspectum*) was found dead on the road between El Río and Sahauripa at the conclusion of

our trip.

In general, we encountered few snakes during our surveys. One neonate black-headed garter snake (*Thamnophis cyrtopsis*) was found at La Cienega and surprisingly was not found at any of the other aquatic sites surveyed. We observed the semi-fossorial Yaqui black-headed snake (*Tantilla yaquia*) in the uplands above Los Pavos camp. This was a fortuitous observation as this species is rarely active on the surface during daylight hours.

Indigo snakes (*Drymarchon corais*), another target species, were the most abundant snake encountered. We observed five individuals at two sites (including one corpse) and obtained photographic vouchers of 2 individuals. All observations of this snake were made in or near dense riparian strand vegetation (Minckley and Brown 1982b).

Conclusions and future research. Amphibians and reptiles of the lower Río Aros and its tributaries have been little studied and few records exist for the region. In contrast, extensive herpetological surveys in Sonora have occurred south of our study area in the vicinity of Alamos, (Bogert and Oliver 1945, Heringhi 1969, Schwalbe and Lowe 2003) and Yécora, Sonora (Van Devender and Ferguson 2003, Lemos-Espinal *et al.* 2004b, Smith *et al.* 2005a, Smith *et al.* 2005b). Surveys to the west of our study area include those by Langebartel and Smith (1954), Smith *et al.* (2005a), and Smith *et al.* (2005b). Based on University of Arizona Herpetological Collections, considerable effort has been conducted to the north as well. Additionally, the western region of Chihuahua, bordering our study area to the east, has received much attention in the last 50 years (Van Devender and Lowe 1977, Tanner 1985, Tanner 1987, Tanner 1989, Lemos-Espinal *et al.* 2004a, Smith *et al.* 2005a, Smith *et al.* 2005b). These surveys were generally focused in the Madrean evergreen woodlands that occur at higher elevations than our study area.

Our observations of spotted box turtles at two sites along the river and a previous observation from Los Pavos Northern Jaguar Preserve (S. Carrillo Percástegui, unpublished data) extends their known range in Mexico by approximately 90 km east-northeast of the nearest known locality (Myers 1945, G. Ferguson pers. comm.). There is essentially nothing known of the life history of spotted box turtles

(Dodd 2001). The closely related ornate box turtle (*Terrepenne ornata luteola*) in Arizona is primarily active during the monsoon season and detectability can be relatively high during this period in appropriate habitat. Based on our observations, it is likely that spotted box turtles follow a similar monsoon-active pattern. Our study area centered on the Los Pavos Northern Jaguar Preserve could provide an ideal conservation area in which to more closely study the life history of spotted box turtles at or near the northern extent of their range.

Indigo snake observations extend the known range of this species in Sonora by approximately 70 km east-northeast of the nearest known record (below dam at Presa Novillo, C. Schwalbe pers. comm.). Indigo snakes were certainly abundant in our study area and likely occur further north of the study area.

Our herpetological inventory was successful in that we were able to extend the known range of two species and to fill distributional gaps for other species, however, this work does not constitute a complete inventory of the region. There is much work that remains to be done in the Rio Aros region, and future work in the study area will certainly yield many more distributional records and natural history observations of the diverse herpetofauna of east-central Sonora.

Birds

Effort and species richness. We surveyed a total of 184.8 km of river corridor along seven reaches that averaged 26.4 km, and surveyed a total of 22.5 km along 8 tributaries that averaged 2.8 km (Table 1).

We detected 80 species of birds, 71 along the river and 61 along tributaries (Table 6). On average, we detected 23% more species along the river corridor (38.3 ± 3.2 species/segment/day) than along tributaries (29.6 ± 3.0 species/transect). We observed 19 species along the river corridor that were not observed along tributaries and only six were associated with open water. Cliff swallow, happy wren, yellow warbler, and brown-headed cowbird (scientific names in Table 6) were detected only along the river corridor and song sparrows were conspicuously absent. We detected nine species along tributaries that were not detected along the river corridor including Montezuma quail, elegant trogon, sulphur-bellied

flycatcher, and rose-throated becard. We also detected 14 species that were not observed along the river or its tributaries on the road between Sahuaripa and Nátorá on 27 July 2005; 10 were associated with oak or oak-pine woodland and four with grassland or shrubland communities not found along the river corridor (Table 7).

Breeding and Residency Status. We confirmed breeding of 16 species that included observations of occupied nests of eight species, dependent young of seven species, and nest building of one species (Table 8). We obtained evidence of probable breeding of 20 species and possible breeding of 34 species. We considered great egret, mallard, and green kingfisher as non-breeders because they were not observed in breeding habitat. We are unsure of the breeding status of common merganser, phainopepla, and black-headed grosbeak and did not consider them possible nesters. Warbling vireo, western tanager, and painted bunting were migrants that likely winter elsewhere, whereas spotted sandpipers were migrants that may winter in the study area. We considered Lucy's warblers as migrants although we expect they commonly breed in riparian areas until early summer.

Frequency of Occurrence. Frequency of occurrence across each river segment and tributary varied among species (Table 8). Most species occurred at relatively few locations with 51% detected at $\leq 33\%$ of river segments or transects and 30% detected at $\leq 13\%$. The most common species we observed included canyon wren and northern cardinal that were detected along all river segments and tributaries and Sinaloa wren, yellow-breasted chat, and five-striped sparrow were detected along 93% (14 of 15) of river segments. Species that were least frequently detected (one observation each) included gray hawk, peregrine falcon, western wood pewee, common yellowthroat, and painted bunting.

Frequency of occurrence for many species varied between river segments and tributaries. Black-vented oriole and curve-billed thrasher were ≥ 2.5 times more abundant along tributaries than along the river corridor whereas frequency of common black-hawk, tropical kingbird, and northern mockingbird were ≥ 4 times higher along the river corridor. Yellow-billed cuckoo, dusky-capped flycatcher, rufous-capped warbler, and streak-backed and hooded orioles were equally abundant along the river and in tributaries (Table 8).

Notable Observations. We observed 12 singing yellow-green vireos; eight were located on slopes between 630 and 670 m along an approximately 1 km stretch of river corridor 10-river km below Nátora and four, including a pair, were in riparian vegetation in Canyon Los Lobos. Yellow-green vireos along the river were in thornscrub that had well-defined shrub, sub-canopy, and canopy layers dominated by mauto (*Lysiloma microphyllum*), and large emergent kapok (*Ceiba acuminata*) up to 10 m tall. In Canyon Los Lobos, habitat was dominated by dense associations of Mexican ebony (*Havardia mexicana*), mauto, bebelama (*Siderloxyton occidentale*), netleaf hackberry (*Celtis reticulata*), Mexican jumping bean (*Sebastiania bilocularis*), and soapberry (*Sapindus saponaria*).

We flushed five military macaws from a large limestone cliff above the river approximately seven river km below Los Chinos and observed two flying down Los Pavos Canyon. Common black-hawks were common along the uppermost 160 km of the river corridor where we detected an average of 0.25 occupied sites per 1 km of river. We also observed common black-hawk along two tributaries, both of which had flowing water. In total we detected common black-hawks at 42 localities, 10 where they were paired and eight where they were with dependent young. We observed black-vented orioles at four localities in riparian vegetation that always included palms (*Sabal uresana* or *Brahea brandegeei*). We heard a single singing happy wren that was singing along the Aros at the mouth of Arroyo Bonito and on the Yaqui near Batui in very dense riparian thickets near the river. Rufous-capped warblers were locally common in rocky canyons with a combination of thornscrub and riparian vegetation and were also detected in thornscrub on slopes along the river corridor. We flushed an adult bald eagle from a hillside at the mouth of Arroyo Bonito and observed a vocalizing pair of peregrine falcons above the river near the mouth of Los Pavos Canyon. We observed a female painted bunting on 28 July approximately 5-river km below Nátora. We flushed a pair of Montezuma quail attending at least three young ≤ 1 to 3 days old at 760 m approximately two km NNE of Tunapa. We detected these Montezuma quail near an arroyo in an open association of thornscrub and mesquite (*Prosopis velutina*) that included scattered low shrubs, bunchgrasses, and forbs; only four Chihuahua oaks (*Quercus chihuahuensis*) were present nearby otherwise the closest oaks were located >800 m away on north-facing hillsides.

Conclusions. High detectability of birds relative to other taxa enabled us to obtain many observations along the river corridor and in selected tributaries. Although effort was limited to only 8 days, conditions along the river were often calm with high visibility that enabled us to identify most of the species we encountered and to investigate further those of interest. Previous surveys of birds along the portions of the Río Aros and Yaqui that we visited are limited to surveys by Bryan Brown who, along with his associates, made several trips by canoe often during winter and spring between 1981 and 1991, primarily to survey bald eagles (Brown and Warren 1985, Brown 1988, Russell and Monson 1998). Other surveys of birds on the Rios Yaqui and Bavispe region were completed in the 1990's and 2000's (A. Flesch, unpubl. data, R. Mesta, pers. comm.) but these data are not yet available. Given limited accessibility, rarity of summer surveys, and our focus on species of conservation and distributional interest, our efforts produced important information in an area where little data were previously available. For example, we observed 47 species that have not been documented previously along and immediately adjacent to the Río Aros (van Rossem 1945, Russell and Monson 1998); two of these species we considered abundant, 18 common, 12 uncommon, and 15 rare.

Many species we detected are typical summer residents in foothills thornscrub or broadleaf-riparian vegetation, yet others are usually associated with more tropical environments to the south (Howell and Webb 1995, Russell and Monson 1998) and were therefore often found only locally. Yellow-green vireos, although rare, were detected in two locations, one where thornscrub structure was tall and vertically stratified and approached structure similar to that of tropical forest, and another location where riparian vegetation was dense and floristically diverse in the middle and lower stories. Although we expect that yellow-green vireos occur locally further to north in the region near the road between Bacadéhuachi to Tecoriname where Russell and Monson (1998) reported a single locality, individuals along Río Aros are near the northernmost extent of their range. Happy wrens were also rare yet seem to be regular summer residents in areas where riparian thickets are exceptionally dense as indicated by observations of signing birds near Batui each year since 2003 (A. Flesch, unpubl. data). Despite these observations, happy wrens were not detected elsewhere on the Yaqui or Aros, or on the Bavispe below

Granados in August 2004 (A. Flesch, unpubl. data). We observed military macaws within 50 km of the northernmost observations of the species (Russell and Monson 1998) and suspect they regularly occur where large limestone cliffs such as those commonly found in and around the Sierra Los Pavos occur near stands of tropical vegetation. Both the location of our observations and those of others (N. Snyder pers. comm. to R. Williams, and R. Williams pers comm.) suggest macaws are present during the breeding season and likely nest on cliffs in these areas.

Other species we observed were associated with the river itself and some are more common to the north. The bald eagle we observed was in an area where they have nested in the past (Brown 1988); bald eagles along the Río Yaqui are at the southernmost extent of their breeding range in mainland Mexico (Howell and Webb 1995). Common black-hawks regularly hunt from perches above water or while perched or walking along shore (Johnsgard 1990). As such, we regularly flushed them while floating the river, and the speed that we floated reduced the likelihood of double counting. Although some common black hawks we observed might not have nested, many were in pairs with dependent young or near large stick nests, suggesting they had. Despite reports of only seven presumed breeding territories of black hawks along the sections of river we floated (Russell and Monson 1998), our findings suggest a larger breeding population, especially given the fact that many black hawks were possibly attending hard-to-detect dependant young. Given the large size of the lower Río Aros relative to other permanent waterways in Sonora and our observations of 42 occupied sites, this area likely supports both the highest densities and largest overall population of common black-hawks in Sonora. By comparison, only five presumed breeding localities have been identified along the Río Bavispe between Granados and the confluence of the Río Aros (A. Flesch, unpubl. data).

Although we obtained information where little was previously available, our efforts were late in the breeding season for many species and produced almost as many questions as we answered. For example, we failed to detect song sparrows despite presence of what seemed to be habitat along the river corridor, our knowledge of their vocalizations, and their typical close proximity to the river bank, suggesting none were present. By comparison, two singing song sparrows were detected along the Río

Bavispe between Granados and the confluence of the Río Aros in mid-August 2004 (A. Flesch, unpubl. data). Although the breeding status of yellow warblers is unknown in our area (Russell and Monson 1998), we observed several singing males in tall stands of willow (*S. gooddingii* and *S. bonplandiana*) especially along the first 30-river km below Nátorá, suggesting the possibility of nesting. We also observed many non-singing males that may have been migrants. Montezuma quail are usually associated with oak woodland but can occur in other vegetation communities at higher elevations or in semidesert grasslands (Corman and Wise-Gervais 2005); in Sonora they primarily occur between 1,200 and 2,000 m (Russell and Monson 1998). Therefore our observation of recently hatched young at only 760 m in thornscrub was unusual. Additional effort will be required to elucidate these questions.

Mammals

To be most effective, camera traps are generally set for long periods of time (e.g. 4-8 weeks). Because of limited trapping periods (~8 hrs) for the duration of the study we only obtained twelve photographs which consisted of: humans (n=6), a domestic dog (*Canis familiaris*) (n=2), a coyote (*Canis latrans*) (n=1), a mouse (*Peromyscus*) (n=1), and two blank photos.

We sampled small mammals with Sherman traps at three camps (see Table 1, Figure 2), but unfortunately, perhaps due to overnight rain, our trap success was very poor. At Buena Vista camp we caught four individuals: three pocket mice (*Chaetodipus penicillatus*) and one kangaroo rat (*Dipodomys* spp). The kangaroo rat escaped before processing, thus making identification to species impossible. At Los Lobos, we made no captures and at Los Pavos camp, we caught two desert pocket mice (photo 3a).

We observed a coatimundi (*Nasua narica*) along the river on 30 July between Buena Vista and Los Lobos. It was observed on a rocky point along the river and upon noticing our river party moved away from the river corridor. We were able to get a positive visual identification and a photographic voucher (Photo 3b). Additionally, we also observed a coatimundi at Los Lobos canyon, although we could not obtain a photo. Additional sightings of mammals included rock squirrel (*Spermophilus variegatus*), and a cottontail (*Sylvilagus* spp.; Table 9). We recorded large felid (i.e., mountain lion, jaguar) scat and cougar tracks at three sites: Buena Vista, Carrizoso, and los Pavos (Photo 4a, 4b). A large felid scrape was also

recorded at Buena Vista on a tree morning glory (*Ipomoea arborecens*, Table 1), and a mountain lion (*Felis concolor*) skull was found and photographed at this camp. The presence of Neotropical river otter tracks at the confluence of the Río Bonito and Río Aros were significant findings since there are very few published records from the state of Sonora (Gallo-Reynosa 1996). This sighting coincides with previous observations from the area (O. Rosas and B. Long, pers. comm.).

Conclusion. Because of the brevity of the research trip we detected a low number of species and individual mammals in the study region as compared to other taxonomic groups. This by no means implies that the region does not harbor a great diversity of mammal species; certainly more effort is needed in this regard.

Current studies of the mammal fauna in NE Sonora include those on beaver (*Castor canadensis frondator*) in the upper Rio Bavispe (Gallo-Reynosa *et al.* , 2002), carnivores in general (Lorenzana-Piña *et al.* 2004), Jaguars (López-González and Lorenzana-Piña 2001, López-González and Brown 2002), and black bears (*Ursus americanus*) and cougars (Carlos López-González, University of Queretaro). López-González and associates have also studied medium and large mammals in East Central Sonora using camera traps (Lorenzana-Piña *et al.* 2004). Other studies include those by Octavio Rosas and Raul Valdez (New Mexico State University) on large felids in the Aros region. Currently, important conservation projects and research on jaguars is being conducted on the Los Pavos Northern Jaguar Preserve.

In-depth studies of species of biogeographic and conservation interest, such as the neotropical river otter and ocelot are lacking in the study area. Additionally, the population dynamics of collared peccary (*Pecari tajacu*) and white-tailed deer (*Odocoileus virginianus*), both important prey species for large felids, remain unknown in the region. Clearly, more work is needed for a more comprehensive view of the geographic distribution and conservation issues of mammals along the Río Aros and its tributaries. We hope that these initial findings not only add to the body of knowledge about the mammal fauna of the region, but also generate interest for future studies.

CONCLUSIONS

Preliminary surveys of the flora and fauna of the Río Aros region support our prediction that an inventory in the region would identify notable observations and range extensions for multiple taxa. In the limited eight day survey period we identified many such species and report the range extension of three species of plants, four aquatic invertebrates, and two reptiles, with numerous other observations of biogeographic note.

The Río Aros region lies at the border of temperate and subtropical biogeographic provinces, however the majority of notable species we encountered were species with ranges extending to the south, likely due to the fact that the dominant vegetation type in this area is subtropical in origin (Brown 1982). At higher elevations upstream we noted large areas of frost-killed vegetation, suggesting that infrequent frost events likely limit some plant species and associated animals in the region. Whereas frost could be a limiting factor for vegetation, groups such as aquatic invertebrates would be unaffected by brief periods of freezing weather, suggesting that global warming could differentially affect the distributional ranges of different plant and animal groups in the region. Further work should identify the role of rare weather events on the distribution of plants in Aros and Yaqui valleys.

Our findings clearly underscore the need for future work in the region. For logistical reasons we chose to conduct our study during the monsoon season when water flows were elevated. However, research during other seasons would allow a better assessment for certain groups. During the monsoon period many reptiles and amphibians may be more detectable, however, other groups such as aquatic invertebrates and some birds may be harder to detect.

This study contributes significantly to biological knowledge of the Río Aros and upper Río Yaqui region. Not only did we compile preliminary species lists that can be used for future efforts in the region, we also identified range extensions for nine species and gained further biogeographic information for many other notable species, underscoring the importance of this region as an outpost for northern limits of subtropical flora and fauna in North America. This region harbors a labyrinth of canyons, mesas, and mountains that await further, more detailed exploration by naturalists. Should additional efforts be made,

we trust the findings will benefit our knowledge of plant and animal distribution and ecology and be of great reward to those making the effort. We hope this study inspires such work and fosters conservation efforts in the future.

ACKNOWLEDGEMENTS

Support for this project was received from T & E Inc, and without their sponsorship, and that of Tom Wootten this project would not have been possible. Mac Hudson, Amy Schwemm, and Sara Jensen provided additional monetary assistance. Lane Larson and Caimen Enterprises provided logistical support and knowledge of the region. In addition, we appreciate the assistance of Matt Havlick and Kevin, Sue and Shannon Green. We gratefully appreciate the endorsement of Northern Jaguar Project, Naturalia, and the Northern Jaguar Preserve and their permission to conduct fieldwork on the Los Pavos-Northern Jaguar Preserve. Raven Bolas from the University of Arizona herbarium assisted with plant identification. Sara Jensen and Thomas van Devender reviewed earlier drafts of this manuscript and provided valuable suggestions. Cory Jones of Sky Island Alliance assisted with the preparation of Figure 1.

LITERATURE CITED

- Arriaga, L., J.M. Espinoza, C. Aguilar, E. Martínez, L. Gómez, and E. Loa (coords.). 2000a. Regiones Terrestres Prioritarias de México. CONABIO. México.
- Arriaga, L., V. Aquilar, J. Alcocer, R. Jiménez, E. Muñoz, and E. Vázquez (coords.). 2000b. Regiones Hidrológicas Prioritarias de México. CONABIO. México.
- Bojorquez, L., R. Aguirre, and A. Ortega. 1985. Rio Yaqui watershed, northern Mexico: Use and management. p 475-478 *in* R. R. Johnson, C. D. Ziebell, D. R. Patton, P. F. Ffolliott, and R. H. Hamre, eds. Riparian Ecosystems and their management: Reconciling conflicting uses. First North American Riparian Conference. 523 pp.
- Bogan, M.T. 2005. Diversity and community structure of aquatic insects in isolated montane desert streams. Master's thesis. Oregon State University, Corvallis.
- Bogert, C.M., J.A. Oliver. 1945. Preliminary analysis of the herpetofauna of Sonora. Bulletin of the American Museum of Natural History (83):301-425
- Bradley, G. A., P. C. Rosen, M. J. Sredl, T. R. Jones, and J. E. Longcore. 2002. Chytridiomycosis in native Arizona frogs. Journal of Wildlife Diseases 38:206-212.

- Brown, B. T. 1988. Additional bald eagle nesting records from Sonora. *Journal of Raptor Research* 22:30-32.
- Brown, B. T. and P. L. Warren. 1985. Wintering bald eagles along the Río Yaqui, Sonora, Mexico. *Wilson Bulletin* 97:224-226.
- Brown, D. E. (editor). 1985. *Biotic communities: Southwestern United States and Northwestern Mexico*. University of Utah Press, Salt Lake City.
- Brown, D. E. and C. A. López-Gonzalez. 2001. *Borderland Jaguars (Tigres de la Frontera)*. University of Utah Press, Salt Lake City.
- Brown, D. E. and C. H. Lowe. 1980. *Biotic communities of the southwest*. U.S. Department of Agriculture Forest Service GTR-RM-78. Rocky Mountain Range and Experiment Station, Fort Collins, Colorado.
- Childs, J. 1998. *Tracking the Felids of the Borderlands*. Printing Corner Press, El Paso.
- Clarkson, R. W. and J. C. Rorabaugh. 1989. Status of leopard frogs (*Rana pipiens* complex: Ranidae) in Arizona and southeastern California. *Southwestern Naturalist* 34:531-538.
- Corman, T. E. and C. Wise-Gervais, editors. 2005. *Arizona breeding bird atlas*. University of New Mexico Press, Albuquerque, New Mexico.
- Crump, M. L. and N. J. Scott, Jr. 1994. Visual encounter surveys. Pages 84-92 in W. R. Heyer, M. A. Donnelly, R. W. McDiarmid, L. C. Hayek, and M. S. Foster, editors. *Measuring and monitoring biological diversity: Standard methods for amphibians*. Smithsonian Institution Press, Washington D.C.
- Dodd, C.K. 2001. *North American Box Turtles: A Natural History*. University of Oklahoma Press, Norman, Oklahoma.
- Elbroch, M. 2003. *Mammal tracks and Sign: A guide to North American species*. Stackpole Books. Mechanicsberg, PA.
- Ernst, C.H. and R.W. Barbour. 1989. *Turtles of the world*. Smithsonian Institution Press, Washington, D.C.
- Felger, R. S., M. B. Johnson, and M. F. Wilson. 2001. *The trees of Sonora, Mexico*. New York: Oxford University Press.
- Frost, J.S., and J.T. Bagnara. 1976. A new species of leopard frog (*Rana pipiens* complex) from Northwestern Mexico. *Copeia* 1976:332-338.
- Gallo-Reynoso, J. P. 1996. Distribution of the Neotropical River otter (*Lutra longicaudis annectens* Major 1987) in the Rio Yaqui, Sonora Mexico. *IUCN Otter Spec. Group. Bull* 13: 27-31.
- Gallo-Reynoso, J. P., G. Suárez-Gracida, H. Cabrera-Santiago, E. Coria-Galindo, J. Egado-Villarreal, and L. C. Ortiz. 2002. Status of beaver (*Castor canadensis frontator*) in Río Bavispe, Sonora, Mexico. *The Southwestern Naturalist* 47:501-504.

- Gentry, H. S. 1982. Sinaloan deciduous forest. *Desert Plants* 4:73-77
- Hale, S.F., G.M. Ferguson, P.A. Holm, and E.B. Wirt. 1998. Re-survey of selected Tarahumara frog (*Rana tarahumara*) localities in Northern Sonora, Mexico, in May 1998. Report submitted to the Arizona Zoological Society and the Tarahumara Frog Conservation Team.
- Heringhi, H.L. 1969. An ecological survey of the herpetofauna of Alamos, Sonora, Mexico. M.S. Thesis. University of Arizona, Tucson, Arizona.
- Hillis, D.M. and T.P. Wilcox. 2005. Phylogeny of the New World true frogs (*Rana*). *Molecular Phylogenetics and Evolution* 34:299-314.
- Howell, S. N. G. and S. Webb. 1995. A guide to the birds of Mexico and northern Central America. Oxford University Press, Oxford, U. K.
- Jasper, S. K., and G. L. Challet. 2004. A Genus and Species New to the USA, *Macrovatellus mexicanus* Sharp (Coleoptera: Dytiscidae), in South Texas, with Habitat Notes. *Coleopterists Bulletin* 56:142-143.
- Johnsgard, P. A. 1990. Hawks, eagles, and falcons of North America: biology and natural history. Smithsonian Institution Press, Washington D.C.
- Langebartel, D.A. and H.M. Smith. 1954. Summary of the Norris collection of reptiles and amphibians from Sonora, Mexico. *Herpetologica* 10:125-136.
- Lemos-Espinal, J.A., H.M. Smith, D. Chizar. 2004a. Introduction to the amphibians and reptiles of the state of Chihuahua, Mexico. Mexico, D.F.: Comisin Nac. para el Conocimiento y uso de la Biodiversidad. 128 p.
- Lemos-Espinal, J.A., H.M. Smith, D. Hartman, and D. Chizar. 2004b. Selected year 2003 amphibians and turtles from Chihuahua and Sonora, Mexico. *Bulletin of the Chicago Herpetological Society* 39:107-109.
- López-González, C. A., and D. E. Brown. 2002. Distribución y estado de conservación actuales del jaguar en el noreste de México. Pp. 379-391 in R. A. Medellín, E. Equihua, C. L. B. Chetkiewickz, P. G. Crawshaw Jr., A. Rabinowitz, K. H. Redford, J. G. Robinson, E. W. Anderson, and A. Taber, comps. *El jaguar en el Nuevo Milenio*. Fondo de Cultura Económica, Universidad Nacional Autónoma de México, Wildlife Conservations Society, México City, México.
- López-González, C. A. and G. Lorenzana-Piña. 2001. Sonoran jaguars: conserving border-lands cats. *Wild Earth* 11: 60-65.
- Lorenzana-Piña, G, R. Castillo and C. A. Lopez-Gonzalez. 2004. Distribution, Habitat Association, and Activity Patterns of Medium and Large Sized mammals of Sonora, Mexico. *Natural Areas Journal*. 24: 354-357
- Lynam, A. J. 2002. Metodos de trabajo de campo para definir y proteger poblaciones dde gatos grandes: los tigres indochinos como un estudio de caso. pp 55-71 in R. A. Medellín, C. Equihua, C. L. B. Chetkiewickz P. G. Crashaw Jr., A. Rabinowitz, K. H. Redford, J. G. Robinson, E. W. Anderson and

- A. Taber comps. El jaguar en el nuevo milenio. Fondo de Cultura Economica. Universidad Nacional Autonoma de Mexico. Wildlife Conservation Society, Mexico City, Mexico.
- Lytle, D. A. 2003. Reconstructing long-term flood regimes with rainfall data: effects of flood timing on caddisfly populations. *The Southwestern Naturalist* 48:36-42.
- McCune, B. and J. B. Grace. 2002. *Analysis of Ecological Communities*. MjM Software Design. Glenden Beach, OR.
- McCune, B. and M. J. Mefford. 1999. *PC-ORD. Multivariate Analysis of Ecological Data. Version 4.0*. MjM Software, Glenden Beach, Oregon, USA.
- MacArthur, R. H. 1972. *Geographical ecology: patterns in the distribution of species*. Harper and Row, New York.
- Marshall, J. T. Jr. 1957. Birds of pine-oak woodland in southern Arizona and adjacent Mexico. *Pacific Coast Avifauna* 32:1-125.
- Martin, P. S., D. Yetman, M. Fishbein, P. Jenkins, T. R. van Devender, and R. Wilson, editors. 1998. *Gentry's Río Mayo plants: the tropical deciduous forest and environs of northwest Mexico*. University of Arizona Press, Tucson, Arizona.
- Mickley, W.L. and D.E. Brown. 1982a. Sinaloan riparian evergreen forest and woodland. *Desert Plants* 4:575
- Mickley, W.L. and D.E. Brown. 1982b. Warm-temperate interior strands. *Desert Plants* 4:265-267.
- Mielke, P. W., Jr, and K. J. Berry. 2001. *Permutation Methods: A distance function approach*. Springer, Berlin.
- Myers, G.S. 1945. A third record of the Sonoran box turtle. *Copeia* 1945:172.
- Platz, J.E. and J.S. Frost. 1984. *Rana yavapaiensis*, a new species of leopard frog (*Rana pipiens* complex). *Copeia* 1979: 383-390.
- Platz, J. E., and J. S. Mecham. 1979. *Rana chiricahuensis*, a new species of leopard frog (*Rana pipiens* Complex) from Arizona. *Copeia* 1979: 383-390.
- Revenga, C., S. Murray, J. Abramovitz, and A. Hammond, 1998. *Watersheds of the World: Ecological Value and Vulnerability*. Washington, DC: World Resources Institute.
- Robichaux, R. H. and D. Yetman, editors. 2000. *The tropical deciduous forest of Alamos: Biodiversity of a threatened ecosystem in Mexico*. University of Arizona Press, Tucson, Arizona.
- Russell, S. M. and G. Monson. 1998. *The birds of Sonora*. University of Arizona Press, Tucson, Arizona.
- Schwalbe, C. R., and C. H. Lowe. 2003. Amphibians and reptiles of the Sierra de Alamos. Pp. 172-199 in R. H. Robichaux and D. A. Yetman (eds.) *The Tropical Deciduous Forest of Alamos*, University of Arizona Press, Tucson.

- Smith, H.M., J.A. Lemos-Espinal, and D.Chizar. 2005a. 2004 snakes from Sonora, Chihuahua and Coahuila, Mexico. *Bulletin of the Chicago Herpetological Society* 40:66-70.
- Smith, H.M., J.A. Lemos-Espinal, and D.Chizar. 2005b. 2004 amphibians from Sonora, Chihuahua and Coahuila, Mexico. *Bulletin of the Chicago Herpetological Society* 40:45-51.
- Stebbins, R.C. 2003. *Western reptiles and amphibians*. Houghton Mifflin Company, Boston.
- Tanner, W.W. 1985. Snakes of Western Chihuahua. *Great Basin Naturalist* 45:615-676.
- Tanner, W.W. 1989. Amphibians of Western Chihuahua. *Great Basin Naturalist*. 49:38-70.
- Tanner, W.W. 1987. Lizards and turtles of Western Chihuahua. *Great Basin Naturalist* 47:383-421.
- Taylor, W. P. 1934. Significance of extreme and intermittent conditions in distribution of species and management of natural resources, with a restatement of Leiberg's law of the minimum. *Ecology* 15:374-379.
- Truxal, F. S. 1953. A revision of the genus *Buenoa* (Hemiptera, Notonectidae). *University of Kansas Science Bulletin* 35:1353-1523.
- Truxal, F. S. and A. S. Menke. 1966. New Distribution Data for *Martarega*, *Buenoa* and *Abedus*, including the first record of the genus *Martarega* in the United States (Hemiptera: Notonectidae: Belostomatidae). *Los Angeles County Museum Contributions in Science* 106:1-6
- Turner, R. M., J. E. Bowers, and T. L. Burgess. 1995. *Sonoran Desert plants: an ecological atlas*. University of Arizona Press, Tucson, Arizona.
- Van Devender, T. R. and C. H. Lowe, Jr. 1977. Amphibians and reptiles of Yepómera, Chihuahua, Mexico. *Journal of Herpetology*. 11:41-50.
- Van Devender, T.R. and G.M. Ferguson. 2003. Research in the Sierra Madre Occidental of Eastern Sonora, Mexico: Noteworthy Animals in the Yecora Area. http://www.desertmuseum.org/programs/yecora_fauna.html
- Van Rossem, A. J. 1945. A distributional survey of the birds of Sonora, Mexico. *Occasional Papers Museum of Zoology, Louisiana State University* 21:1-379.
- Williams, D. D., and H. B. Hynes. 1976. The recolonization mechanisms of stream benthos. *Oikos* 27:265-272.

Figure 2. Detailed map of the study region. River sections between red slashes (campsites) are labeled with capital letters. Campsites are designated with roman numerals and take the name of the nearest geographical placename. Surveyed tributaries are designated with Arabic numbers. An inset map shows the location of the study area within Mexico.




Figure 3. Non-metric multidimensional scaling ordination graph of tributary and mainstem samples, with significantly associated physical variable (flow) overlain.




Table 1. Summary of river reaches (Ríos Aros and Yaqui), tributaries, and campsites surveyed.

River reach	code	Date	(km)	Tributary	Map code	Date	Distance	Plant	Bird	Invert.	Herps	Mammals
Natora to Tunapa	A	7/27-28	21.0	Los Chinos	1	7/28-29	3.3	X	X		X	X
Tunapa to Buena Vista	B	7/29	33.0	La Cienega	2	7/29	2.5	X	X		X	
Buena Vista to Los Lobos	C	7/30	29.6	Unnamed at	3	7/28-29	1.5			X	X	X
Los Lobos to El Carrizoso	D	7/31	29.7	Tunapa camp								
Carrizoso to Los Pavos	E	8/1	11.4	El Placer	4	7/30	3.0	X	X	X		
Los Pavos to Los Alisos	F	8/2	34.8	Chino Gordo	5	7/30	1.8	X	X		X	X
Los Alisos to El Rio	G	8/3	25.3	Río Bonito	6	7/30	0.5			X	X	X
				Los Lobos	7	7/31	2.6	X	X	X	X	X
River camp	code	Date		Carrizoso	8	8/1	2.5	X	X	X	X	X
Natora	I	7/27-28		El Toro	9	8/2	1.0			X	X	
Tunapa	II	7/29-30		Los Pavos	10	8/2	1.4	X	X		X	X
Buena vista	III	7/30-31		Fig-fall	11	8/2	1.0				X	
Los lobos	IV	7/31-8/1		Los Alisos	12	8/3	5.4	X	X	X	X	
El Carrizoso	V	8/1-2										
Los Pavos	VI	8/2-3										
Los Alisos	VII	8/3-4										

Table 2. Sampling locations and abiotic conditions for aquatic invertebrate sampling. All UTM coordinates taken using the NAD 1927 datum.

Sites	Date	N UTM	E UTM	Water°C	pH	Cond (S)	Estimated flow (cfs)
<i>Tributary Canyons</i>							
Carrizoso	1-Aug-05	3266472	679414	28	8	480	0.1
El Placer	30-Jul-05	3239426	702416	23	8	310	3
El Toro	1-Aug-05	*	*	28	*	*	0.01
Chino Gordo	30-Jul-05	3238838	704898	*	*	*	0.01
Los Lobos	31-Jul-05	3249866	692628	25	7.75	590	0.01
Unnamed trib. @ Tunapa camp	29-Jul-05	3217863	718064	*	*	*	0.1
Fig-Waterfall	1-Aug-05	3259176	669758	29	*	*	0
High Elevation Creek b/w Nátora & Sahuaripa	27-Jul-05	3212880	695440	20	*	*	0
<i>Mainstem- Drift Samples</i>							
El Carrizoso camp	31-Jul-05	3266394	679294	25	7.75	60	3200
Los Alisos camp	3-Aug-05	3249780	671174	24.5	7.25	*	4000
<i>Mainstem- Net Samples</i>							
Río Aros at Buena vista camp	30-Jul-05	3238838	704898	26	7.75	60	3200
Río Aros at Río Bonito	30-Jul-05	3250409	698062	27	*	*	3200
Río Aros at Tunapa camp	28-Jul-05	3217683	718064	27	*	*	2500
Río Sahuaripa at road xing to Nátora	27-Jul-05	3210117	674656	*	*	*	1000
Río Yaqui at takeout	3-Aug-05	3231824	667642	25	*	*	3500

* no data taken

Table 3. Species occurrences of aquatic invertebrates by sampling location.

Taxa	Tributary Canyons								Mainstem- Drift		Mainstem- Net				
	Carrizoso	Chino	Fig-Fall	High Elv.	Lobos	Placer	Toro	Tunapa	Alisos	Carrizoso	Bonito	Chinos	Sahuaripa	Takeout	Tunapa
COLEOPTERA															
Dryopidae															
sp.					x										
<i>Dryops arizonensis</i>					x										
<i>Postelichus</i>	x														
Dytiscidae															
<i>Desmopachria dispersa</i>							x								
<i>Desmopachria mexicana</i>	x		x		x		x	x							
<i>Desmopachria portmanni</i>				x	x										
<i>Laccophilus fasciatus</i>	x		x			x	x	x							
<i>Laccophilus horni</i>					x										
<i>Laccophilus mexicanus</i>	x				x										
<i>Laccophilus pictus</i>	x		x	x	x		x	x					x		
<i>Liodessus obscurellus</i>				x										x	
<i>Macrovatellus mexicana</i>	x				x										
<i>Neoclypeodytes</i>	x			x	x			x					x		
<i>Rhantus atricolor</i>				x											
<i>Rhantus calidus</i>					x										
<i>Rhantus gutticollis gutticollis</i>				x											
<i>Stictotarsus aequinoctialis</i>				x	x										
<i>Stictotarsus roffi</i>	x				x								x		
<i>Stictotarsus striatellus</i>					x										
<i>Thermonectus marmoratus</i>			x	x	x		x	x							
Elmidae										x		x	x		
Gyrinidae															
<i>Dineutus sublineatus</i>	x			x				x							
<i>Gyretes torosus</i>	x														
<i>Gyrinus plicifer</i>				x											
Haliplidae															
<i>Peltodytes dispersus</i>	x				x			x							
Helophoridae															
<i>Helophorus</i>	x														
Heteroceridae															
Hydraenidae															
<i>Hydraena</i>				x	x										
<i>Ochthebius discretus</i>					x										
Hydrochidae															
<i>Hydrochus</i>	x				x										

Table 3. (continued). Species occurrences of aquatic invertebrates by sampling location.

Taxa	Tributary Canyons								Mainstem- Drift		Mainstem- Net				
	Carrizoso	Chino	Fig-Fall	High Elv.	Lobos	Placer	Toro	Tunapa	Alisos	Carrizoso	Bonito	Chinos	Sahuaripa	Takeout	Tunapa
COLEOPTERA															
Hydrophilidae															
<i>Anacaena limbata</i>					x										
<i>Berosus blechrus</i>				x	x		x								
<i>Berosus miles</i>			x												
<i>Berosus moerens</i>				x	x	x								x	
<i>Berosus notapeltatus</i>				x	x										
<i>Berosus</i> nr. <i>salvini</i>	x														
<i>Berosus</i> nr. <i>stylifer</i>														x	
<i>Berosus rugulosus</i>	x		x		x	x	x	x	x					x	
<i>Berosus salvini</i>				x											
<i>Berosus</i> sp.	x							x							
<i>Chaetarthria</i> sp. 1					x										
<i>Chaetarthria</i> sp. 2					x										
<i>Enochrus</i>	x		x		x		x								
<i>Helochares normatus</i>	x		x		x		x								
<i>Hydrophilus triangularis</i>	x													x	
<i>Paracymus</i>	x							x							
<i>Tropisternus ellipticus</i>	x		x	x				x						x	
<i>Tropisternus lateralis</i>	x							x						x	
Scirtidae															
<i>Cyphon</i>					x										
Staphlynidae															
				x	x										
DIPTERA															
Ceratopogonidae															
					x										
Chironomidae															
	x				x					x				x	
Culicidae															
	x				x										
Simuliidae															
Stratiomyidae															
<i>Oxycera</i>					x										
<i>Stratiomys</i>			x											x	
Tabanidae															
<i>Tabanus</i>														x	
EPHEMEROPTERA															
Baetidae															
<i>Callibaetis</i>	x			x	x	x								x	
Caenidae															
<i>Caenis</i>														x	
Leptophlebiidae															
<i>Traverella</i>										x			x		x

Table 3. (continued). Species occurrences of aquatic invertebrates by sampling location.

Taxa	Tributary Canyons								Mainstem- Drift		Mainstem- Net				
	Carrizoso	Chino	Fig-Fall	High Elv.	Lobos	Placer	Toro	Tunapa	Alisos	Carrizoso	Bonito	Chinos	Sahuaripa	Takeout	Tunapa
LEPIDOPTERA															
Pyrilidae									x						
MEGALOPTERA															
Corydalidae															
<i>Corydalus</i>											x	x			x
ODONATA															
Aeshnidae															
<i>Aeshna</i>					x										
Calopterygidae															
<i>Hetaerina americana</i>	x														
Coenagrionidae															
sp.					x							x			
<i>Argia</i>	x						x	x							
Libellulidae	x				x										
TRICHOPTERA															
Calamoceratidae															
<i>Phylloicus aeneus</i>					x										
Hydropsychidae															
<i>Smicridea</i>									x	x					
Odontoceridae															
<i>Marilia flexuosa</i>							x	x							

Table 4. Amphibian and reptile species detections by site.

	<u>Site Name / Tributary Canyon</u>														
	High Elev	Natora	Tunapa	La Cienega	Buena Vista	Chino Gordo	Arroyo Bonito	Los Lobos	Carrizoso	El Toro	Los Pavos	Fig Fall	Los Alisos	El Rio	
Salamanders															
<i>Ambystoma rosaceum</i>	X														
Frogs and toads															
<i>Gastrophryne olivacea</i>				X					X						
<i>Hyla arenicolor</i>						X			X			X	X		
<i>Bufo alvarius</i>									X	X					
<i>Bufo mazatlanensis</i>		X	X		X										
<i>Bufo punctatus</i>			X			X		X				X			
<i>Rana tarahumarae</i>				X											
<i>Rana cf. yavapaiensis</i>	X						X	X	X				X	X	
Turtles															
<i>Trachemys scripta</i> subsp.					X							X			
<i>Terrepenne nelsoni</i>					X		X								
<i>Kinosternon sonoriense</i>	X		X	X		X	X								
Lizards															
<i>Heloderma suspectum</i>														X	
<i>Callisaurus draconoides</i>											X			X	
<i>Holbrookia maculata</i>			X												
<i>Sceloporus clarkii</i>	X				X	X			X?	X					
<i>Urosaurus ornatus</i>						X					X	X	X		
<i>Eumeces obsoletus</i>															
<i>Aspidoscelis spp.</i>									X						
Snakes															
<i>Drymarchon corais</i>			X			X			X						
<i>Tantilla yaqui a.</i>											X				
<i>Thamnophis cyrtopsis</i>				X											

Table 5. Species list for reptiles and amphibians. Common names and Latin nomenclature given.

Scientific Classification and Name	Common Name
Caudata	Salamanders
Ambystomatidae	Mole Salamanders
<i>Ambystoma rosaceum</i>	Tarahumara Salamander
Anura	Frog and Toads
Microhylidae	Narrow-mouthed Toads
<i>Gastrophryne olivacea mazatlanensis</i>	Sinaloan narrow-mouthed toad
Hylidae	Treefrogs
<i>Hyla arenicolor</i>	Canyon treefrog
Bufonidae	True Toads
<i>Bufo alvarius</i>	Sonoran Desert toad
<i>Bufo mazatlanensis</i>	Mazatlán toad
<i>Bufo punctatus</i>	Red-spotted toad
Ranidae	True frogs
<i>Rana tarahumarae</i>	Tarahumara frog
<i>Rana c.f. yavapaiensis</i>	Lowland leopard frog
Testudines	Turtles
Emydidae	Box and Water turtles
<i>Trachemys scripta</i> subsp.	Pond slider turtle
<i>Terrepene nelsoni</i>	Spotted box turtle
Kinosternidae	Mud turtles
<i>Kinosternon sonoriense</i>	Sonoran mud turtle
Squamata	Lizards
Helodermatidae	Beaded lizards
<i>Heloderma suspectum</i>	Gila monster
Phrynosomatidae	
<i>Callisaurus draconoides</i>	Zebra tailed lizard
<i>Holbrookia maculata</i>	Lesser earless lizard
<i>Sceloporus clarkii</i>	Clark's spiny lizard
<i>Urosaurus ornatus</i>	Tree lizard
Scincidae	Skinks
<i>Eumeces obsoletus</i>	Great Plains skink
Teiidae	Whiptails
<i>Aspidoscelis spp.</i>	Whiptail
Squamata	Snakes
Colubridae	
<i>Drymarchon corais</i>	Indigo snake
<i>Tantilla yaquia</i>	Yaqui black-headed snake
<i>Thamnophis cyrtopsis</i>	Black-headed garter snake

Table 6. Relative abundance and breeding status of bird species ($n = 80$) detected along the Ríos Yaqui and Aros and adjacent tributaries.

Common name	Scientific name	Abundance	Breeding evidence	Comments
Great Blue Heron	<i>Ardea herodias</i>	Abundant	Present in suitable habitat	As many as 70 to 100 per day
Great Egret	<i>Egretta alba</i>	Uncommon	Not in breeding habitat	
Mallard (Mexican variety)	<i>Anus platyrhynchos</i>	Rare	Not in breeding habitat	
Common Merganser	<i>Mergus merganser</i>	Uncommon	Paired, breeding status	Mainly females in groups of 8 or fewer
Black Vulture	<i>Coragyps atratus</i>	Common		
Turkey Vulture	<i>Cathartes aura</i>	Common		
Bald Eagle	<i>Haliaeetus leucocephalus</i>	Rare	At past nesting location	Single adult flushed at mouth of Arroyo Bonito after nesting
Common Black-Hawk	<i>Buteogallus anthracinus</i>	Common	Dependent young	Forty-two sites, 9 w/ dependent young, 10 w/ pairs
Gray Hawk	<i>Buteo nitidus</i>	Rare	Calling territorially	One detection at takeout
Red-tailed Hawk	<i>Buteo jamaicensis</i>	Uncommon	Dependent young	
Peregrine Falcon	<i>Falco peregrine</i>	Rare	Paired and calling	Observed only at mouth of Los Pavos Canyon
Montezuma Quail	<i>Cyrtonyx montezumae</i>	Rare	Dependent young	Pair w/ yg <3 days old in thornscrub w/ few oaks
Elegant Quail	<i>Callipepla douglasii</i>	Common	Singing, paired	
Spotted Sandpiper	<i>Actitis macularia</i>	Common		As many as 14 per day
White-winged Dove	<i>Zenaida asiatica</i>	Common	Paired	
Mourning Dove	<i>Zenaida macroura</i>	Uncommon	Nest w/ 2 eggs	
Inca Dove	<i>Columbina inca</i>	Rare	Singing, paired	
Common Ground-Dove	<i>Columbina passerina</i>	Common	Nest w/ eggs and young	
White-tipped Dove	<i>Leptotila verreauxi</i>	Uncommon	Singing	Found throughout in dense thornscrub
Military Macaw	<i>Ara militaris</i>	Rare	Paired	Group of 2 and 5 near large limestone cliffs
Yellow-billed Cuckoo	<i>Coccyzus americanus</i>	Common	Nest w/ 4 eggs	In riparian groves and thornscrub
Greater Roadrunner	<i>Geococcyx californianus</i>	Uncommon		
Great Horned Owl	<i>Bubo virginianus</i>	Uncommon		Using rock alcoves along River
Broad-billed Hummingbird	<i>Cynanthus latirostris</i>	Common		
Violet-crowned Hummingbird	<i>Amazilia violiceps</i>	Common		
Elegant Trogon	<i>Trogon elegans</i>	Uncommon	Calling territorially	In tall thornscrub and riparian trees in side canyons
Green Kingfisher	<i>Chloroceryle americana</i>	Rare		Not along river and only in wet side canyons
Gila Woodpecker	<i>Melanerpes uropygialis</i>	Common	Dependent young, pairs	

Table 6. Relative abundance and breeding status of bird species ($n = 80$) detected along the Ríos Yaqui and Aros and adjacent tributaries.

Common name	Scientific name	Abundance	Breeding evidence	Comments
Ladder-backed Woodpecker	<i>Picoides scalaris</i>	Common	Paired	
Northern Beardless-Tyrannulet	<i>Camptostoma imberbe</i>	Rare	Singing	
Western Wood-Pewee	<i>Contopus sordidulus</i>	Rare	Singing	Observed only in Sycamores in El Placer Canyon Canyon
Black Phoebe	<i>Sayornis nigricans</i>	Common	Singing, paired	
Vermilion Flycatcher	<i>Pyrocephalus rubinus</i>	Rare	Dependent young	
Dusky-capped Flycatcher	<i>Myiarchus tuberculifer</i>	Common	Singing, paired	In thornscrub on slopes and riparian bottoms
Ash-throated Flycatcher	<i>Myiarchus cinerascens</i>	Uncommon		At lower elevations in more open arid thornscrub
Nutting's Flycatcher	<i>Myiarchus nuttingi</i>	Rare		Detected only once in tall thornscrub
Brown-crested Flycatcher	<i>Myiarchus tyrannulus</i>	Uncommon		Restricted to groves of large chinios and willows
Sulphur-bellied Flycatcher	<i>Myiodynastes luteiventris</i>	Rare	Dependent young	2 family groups in Sycamores in Los Lobos Canyon
Tropical Kingbird	<i>Tyrannus melancholicus</i>	Common	Singing, paired	In riparian groves along Río
Cassin's Kingbird	<i>Tyrannus vociferans</i>	Uncommon		
Thick-billed Kingbird	<i>Tyrannus crassirostris</i>	Uncommon		Single birds in 3 side canyons
Western Kingbird	<i>Tyrannus verticalis</i>	Uncommon		
Rose-throated Becard	<i>Pachyramphus aglaiae</i>	Rare	Nest building, copulating	Found mainly in Sycamores in side canyons
Bell's Vireo	<i>Vireo bellii</i>	Common	Singing	
Warbling Vireo	<i>Vireo gilvus</i>	Rare		Few migrants in riparian trees
Yellow-green Vireo	<i>Vireo flavoviridis</i>	Rare	Singing, paired	8 singing along 2 km of river above Tunapa, 4 in Canyon Los
Common Raven	<i>Corvus corax</i>	Common	Used nest	
Cliff Swallow	<i>Hirundo pyrrhonota</i>	Uncommon	Occupied and used nests	Found at only 3 to 5 sites
Verdin	<i>Auriparus flaviceps</i>	Common	Used nests, paired	
Cactus Wren	<i>Campylorhynchus</i>	Common	Used nests, paired	
Rock Wren	<i>Salpinctes obsoletus</i>	Uncommon	Singing	
Canyon Wren	<i>Catherpes mexicanus</i>	Common	Singing, paired	
Sinaloa Wren	<i>Thryothorus sinaloa</i>	Common	Singing, used nests	Throughout especially in riparian woodland adj
Happy Wren	<i>Thryothorus felix</i>	Rare	Singing	Observed at mouth of A. Bonito and Batui
Black-capped Gnatcatcher	<i>Polioptila nigriceps</i>	Common	Singing, paired	
Northern Mockingbird	<i>Mimus polyglottos</i>	Uncommon		Mainly in lower section of Río
Curved-billed Thrasher	<i>Toxostoma curvirostre</i>	Rare	Calling	

Table 6. Relative abundance and breeding status of bird species ($n = 80$) detected along the Ríos Yaqui and Aros and adjacent tributaries.

Common name	Scientific name	Abundance	Breeding evidence	Comments
Phainopepla	<i>Phainopepla nitens</i>	Uncommon		
Lucy's Warbler	<i>Vermivora luciae</i>	Uncommon		Migrants in riparian vegetation
Yellow Warbler	<i>Dendroica petechia</i>	Common	Singing, others not singing	Restricted mainly to willow groves along Río
Common Yellowthroat	<i>Geothlypis trichas</i>	Rare	Singing	Detected only near Nátorá
Rufous-capped Warbler	<i>Basileuterus rufifrons</i>	Uncommon	Singing	More common in side canyons
Yellow-breasted Chat	<i>Icteria virens</i>	Abundant	Nest w/ 4 eggs	
Summer Tanager	<i>Piranga rubra</i>	Common	Singing, pairs	Often in willows along Río and adjacent thornscrub
Western Tanager	<i>Piranga ludoviciana</i>	Rare		
Canyon Towhee	<i>Pipilo fuscus</i>	Common	Singing, paired	
Rufous-winged Sparrow	<i>Aimophila carpalis</i>	Uncommon	Singing	
Five-striped Sparrow	<i>Amphispiza quinquestriata</i>	Abundant	Dep. young, nest w/ 4 eggs	Singing throughout thornscrub
Black-headed Grosbeak	<i>Pheucticus melanocephalus</i>	Common	Singing	No sign of nesting
Northern Cardinal	<i>Cardinalis cardinalis</i>	Uncommon	Singing, paired	
Blue Grosbeak	<i>Guiraca caerulea</i>	Common	Singing, paired	
Varied Bunting	<i>Passerina versicolor</i>	Common	Singing, paired	
Painted Bunting	<i>Passerina ciris</i>	Rare		1 female approx. 10 km below Nátorá
Bronzed Cowbird	<i>Molothrus aeneus</i>	Uncommon		
Brown-headed Cowbird	<i>Molothrus ater</i>	Uncommon		
Black-vented Oriole	<i>Icterus wagleri</i>	Rare	Paired	9 birds near Buena Vista and in 3 large side canyons, often in
Hooded Oriole	<i>Icterus cucullatus</i>	Uncommon	Nests w/ young	As with previous species often near palms
Streak-backed Oriole	<i>Icterus pustulatus</i>	Uncommon	Occupied and used nests	Mainly in chinós along large side canyons and Río
House Finch	<i>Carpodacus mexicanus</i>	Common	Singing, paired	
Lesser Goldfinch	<i>Carduelis psaltria</i>	Uncommon		

Table 7. Birds species ($n = 14$) detected while traveling the road from Sahuaripa to Nátora (100 km) through Sinaloan thornscrub and Madrean evergreen woodland that were not detected along the Río Aros or Yaqui, Sonora, Mexico 27 July 2005.

Common name	Scientific name
Acorn Woodpecker	<i>Melanerpes formicivorus</i>
Arizona Woodpecker	<i>Picoides arizonae</i>
Northern Flicker	<i>Colaptes auratus</i>
White-striped Woodcreeper	<i>Lepidocolaptes leucogaster</i>
Greater Pewee	<i>Contopus pertinax</i>
Loggerhead Shrike	<i>Lanius ludovicianus</i>
Mexican Jay	<i>Aphelocoma ultramarina</i>
Bridled Titmouse	<i>Parus wollweberi</i>
Bushtit	<i>Psaltriparus minimus</i>
White-breasted Nuthatch	<i>Sitta carolinensis</i>
Eastern Bluebird	<i>Sialia sialis</i>
Hepatic Tanager	<i>Piranga flava</i>
Cassin's Sparrow	<i>Aimophila cassinii</i>
Rufous-crowned Sparrow	<i>Aimophila ruficeps</i>

Table 8. Distribution and abundance of birds ($n = 80$) detected along the Rios Yaqui and Aros and adjacent tributaries from Natora to Batacomachi Sonora, Mexico 27 July to 3 August 2005. Numbers indicate singles, singing males, or pairs, x indicates common species for which abundance was not estimated. Kilometers indicate distance floated along river or distance walked along tributaries. C = colony, p = pair, y = young, NE = nest with eggs, UN = used nest, ON = occupied nest. For example a 9,2p1y indicates we observed the species at 9 locations and at 2 there was a pair and at 1 there was dependant young.

Common name	River							Tributary							
	Natora to Tunapa	Tunapa to Buena Vista	Buena Vista to Los Lobos	Los Lobos to Carrizoso	Carrizoso to Los Pavos	Los Pavos o Los Alisos	Los Alisos to El Rio	Los Chinos	La Cienega	El Placer	Chino Gordo	Los Lobos	Carrizoso	Los Pavos	Los Alisos
Map reference code	A	B	C	D	E	F	G	1	2	3	4	5	6	7	8
Kilometers surveyed	21	33	29.6	29.7	11.4	34.8	25.3	3.3	2.5	3.0	1.8	2.6	2.5	1.4	5.4
Date	7/27-28	7/29	7/30	7/31	8/1	8/2	8/3	7/28-29	7/29	7/30	7/30	7/31	8/1	8/2	8/3
Great Blue Heron	15	50+	70+	50+	x	x	x								
Great Egret															
Mallard (Mexican variety)															
Common Merganser		3	15	5		2									
Black Vulture	x	x	x	x		x	x								
Turkey Vulture	x	x	x	x		x	x					x	x		1
Bald Eagle			1												
Common Black-Hawk	7	6,3py	7,2py	7,2p1y	3,1p	9,2p1y	1			1,1y	1				
Gray Hawk							1								
Red-tailed Hawk		1	1	1	1	3		1	1	1			3		2, 1y
Peregrine Falcon						1p									
Montezuma Quail								6, 3y							
Elegant Quail	3	x	x	x	x	x	x	x	x			1		3	x
Spotted Sandpiper	8	3	4	4	5	14	x								
White-winged Dove	x	x	x	x	x	x	x	x	x	x		20	x	x	x
Mourning Dove	x	x						x		x	x				

	<i>Natura to Tunapa</i>	<i>Tunapa to Buena Vista</i>	<i>Buena Vista to Los Lobos</i>	<i>Los Lobos to Carrizoso</i>	<i>Carrizoso to Los Pavos</i>	<i>Los Pavos o Los Alisos</i>	<i>Los Alisos to El Rio</i>	<i>Los Chinos</i>	<i>La Cienega</i>	<i>El Placer</i>	<i>Chino Gordo</i>	<i>Los Lobos</i>	<i>Carrizoso</i>	<i>Los Pavos</i>	<i>Los Alisos</i>
Inca Dove	x														
Common Ground-Dove	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
White-tipped Dove		2	5	x	x	3	1	5	3	2	5	6	1	3	
Military Macaw				5									1p		
Yellow-billed Cuckoo	4	x	x	x		8	1		3	2	1	3,10N		2	
Greater Roadrunner	1					2		1						2	
Great Horned Owl		2		1					1						
Broad-billed Hummingbird	2	1	x	x	x	x	x	1	x	1		3	2	x	
Violet-crowned Hummingbird	4					1						4	1	2	
Elegant Trogon								1			2			5	
Green Kingfisher		2												2	
Gila Woodpecker	x	x	x	x	x	x	x	1			2,2y	x	1	x	x
Ladder-backed Woodpecker			x	x	x	x	x	1				x	1	x	x
Northern Beardless-Tyrannulet	2	x	1					x	1			2	x	1	
Western Wood-Pewee										1					
Black Phoebe	x	x	x	x	x	x		x	x	x		x			x
Vermilion Flycatcher	x	x	x	x		x	x	x		x				x	x
Dusky-capped Flycatcher		2	2	x		x	x	6	x	4		5	8	1	x
Ash-throated Flycatcher						x	x	1			1			1	x
Nutting's Flycatcher															1
Brown-crested Flycatcher	2		1			1		1		1		2	1	1	
Sulphur-bellied Flycatcher												15,10y			
Tropical Kingbird	20	x	x	x	x	x	5	2			1				
Cassin's Kingbird	2														

	<i>Matorra to Tunapa</i>	<i>Tunapa to Buena Vista</i>	<i>Buena Vista to Los Lobos</i>	<i>Los Lobos to Carrizoso</i>	<i>Carrizoso to Los Pavos</i>	<i>Los Pavos o Los Alisos</i>	<i>Los Alisos to El Rio</i>	<i>Los Chinos</i>	<i>La Cienega</i>	<i>El Placer</i>	<i>Chino Gordo</i>	<i>Los Lobos</i>	<i>Carrizoso</i>	<i>Los Pavos</i>	<i>Los Alisos</i>
Thick-billed Kingbird										1			1		
Western Kingbird						5									
Rose-throated Becard									1UN		3,1ON				
Bell's Vireo	12	x	2	x	x	x	x	x	2			x	x	x	
Warbling Vireo					1							1		1	
Yellow-green Vireo	8										4				
Common Raven		x	x									x			x
Cliff Swallow		2C	1				1C								
Verdin	x							x	2					x	x
Cactus Wren				x		x	x	x			x		x	x	x
Rock Wren	x				x								x		
Canyon Wren	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Sinaloa Wren	40+	20+	x	x	x	x	x	x	x	10	3	x	x		2
Happy Wren			1				1								
Black-capped Gnatcatcher	2	3			2	x		x		x		x	x	x	6
Northern Mockingbird	1	1		2		x									2
Curved-billed Thrasher						1	1	2		1		1	1	x	1
Phainopepla													1		
Lucy's Warbler	3							2							
Yellow Warbler	7	x	2	x			x								
Common Yellowthroat	1														
Rufous-capped Warbler		1		2	3					5		5	10		1
Yellow-breasted Chat	x	x	x	x	x	x	x	x	x	x	x		x	x	x
Summer Tanager	5	10	2	x		x	x	x		4		2			x

	<i>Natora to Tunapa</i>	<i>Tunapa to Buena Vista</i>	<i>Buena Vista to Los Lobos</i>	<i>Los Lobos to Carrizoso</i>	<i>Carrizoso to Los Pavos</i>	<i>Los Pavos o Los Alisos</i>	<i>Los Alisos to El Rio</i>	<i>Los Chinos</i>	<i>La Cienega</i>	<i>El Placer</i>	<i>Chino Gordo</i>	<i>Los Lobos</i>	<i>Carrizoso</i>	<i>Los Pavos</i>	<i>Los Alisos</i>
Western Tanager	1	1													
Canyon Towhee	2	x						x		x				x	1
Rufous-winged Sparrow	x					x	x	x						x	
Five-striped Sparrow	3	x	x	x	x	x	x	x	1NE	x		x	x	x	x
Black-headed Grosbeak	1	x				x	x	2				1	1		1
Northern Cardinal	x	x	x	x	x	x	x	x	x	x	x	1	x	x	x
Blue Grosbeak	x	x	x		x	x	x	x	x	x		x	x	x	x
Varied Bunting	x	x	x		x	x	x	x	x	x		x	x	x	1
Painted Bunting	1														
Bronzed Cowbird	2	x	x							x					
Brown-headed Cowbird		x	x	x											
Black-vented Oriole		1										3	3		1
Hooded Oriole	7	x	1	x		x	x	3	x	2		1	3	1	10N
Streak-backed Oriole	2UN	3UN	3UN					10N		2UN		2UN			
House Finch	x		x							x		x	x	1	x
Lesser Goldfinch								2							
Species richness - transect	47	45	40	34	25	40	37	37	19	33	12	32	34	29	41
- river and				71								61			
- region								80							

Table 9. Mammal species detections by site. Codes indicate detection type: SC=scat, SP=scrape, TR=track, P=picture, C=capture, SI=sighting.

Species	Common name English/ Nombre comun español	Site Name/Tributary Canyon					
		Unnamed at Natora	Funapa cam Funapa	Buena Vista	Arroyo Bonito	Los Lobos	Carrizoso los Pavos
<i>Odocoileus virginianus</i>	White tail deer/Venado cola blanca		TR	TR		TR	TR
<i>Pecari tajacu</i>	Javelina/Cochi jabalí		TR		SC	TR	TR
<i>Canis latrans</i>	Coyote/Coyote	TR	SC		TR/SC	PI	TR
<i>Urocyon cinereoargenteus</i>	Grey fox/Zorra gris			TR/SC		TR	TR
<i>Puma concolor</i>	Puma/León			SC/SP			TR/SC
<i>Lynx rufus</i>	Bobcat/Gato pochi			SC			TR
<i>Conepatus mesoleucus</i>	Hog-nosed skunk/Zorrillo cadeno						TR
<i>Spilogale putorius</i>	Spotted skunk/Zorrillo manchado		TR				
<i>Bassariscus astutus</i>	Ringtail/Cacomixtle		TR			TR	TR
<i>Procyon lotor</i>	Raccoon/Batepi	TR	TR		TR	TR	TR
<i>Nasua narica</i>	Coatimundi/Chulo		TR		SI	SI	TR
<i>Lontralongicaudis</i>	Neotropical river otter/nutria				TR		
<i>Lepus alleni</i>	Antelope Jack Rabbit/Liebre	SI					
<i>Sylvilagus sp</i>	Cottontail/Conejo		SI				TR
<i>Spermophilus variegatus</i>	Rock squirrel/Ardilla					SI	
<i>Chaetodipus penicillatus</i>	Desert pocket mouse/Raton			CA			CA
<i>Dipodomys sp.</i>	Kangaroo rat/Rata canguro			CA			
<i>Peromyscus sp</i>	Mouse/Ratón de campo		TR?				TR

Photo 1. Tributary, Carrizoso Canyon, (A). Mainstem Río Yaqui at Los Pavos camp (B).



Photo 2. Dobsonfly (Corydalidae: *Corydalus*) adult (A) and larvae (B) from the Río Aros.



Photo 3. *Chaetodipus penicillatus*, desert pocket mouse, captured at Los Lobos Camp (A). Coatimundi (*Nasua narica*) (B).



Photo 4. Mountain lion scat (A) and tracks (B) from Buena Vista



Photo 5. View from above Tunapa camp, looking WSW. Note top-killed vegetation (appearing brown) on slopes above the river in center-left of photo.

